

This list generally follows the order found in the Miller and Brown (1981) *Catalogue/Checklist of the Butterflies of North America* and the supplement to that list by Ferris (1989c), both published as *Memoirs of the Lepidopterists' Society*. Generic name usage is conservative (when opinions vary on how to divide monophyletic groups) and generally follows that used in *A Field Guide to Eastern Butterflies* (Opler 1992, 1998), *A Field Guide to Western Butterflies* (Opler 1999) and the Stanford and Opler (1993) *Atlas to Western USA Butterflies* (updated as Opler et al. 2000). Some changes are made to conform with recent research results (e.g. Emmel 1998) and with Palaearctic and Neotropical publications (see References).

The list that follows includes superfamily, family, and subfamily categories (with tribes for some Lycaenidae), generally in accord with the arrangement presented by de Jong et al. (1996) and Ackery et al. (1999). The Listing order of families, subfamilies, genera, and species does not necessarily imply relationships of taxa to each other. It is the intent of the authors to use the original spelling of species names as described. We would be pleased if anyone would point out where names herein do not conform to their original orthography.

The main purpose of this list is to summarize our current knowledge of the nomenclature of North American butterflies at the species-level (and higher), and to generate interest in current nomenclatural problems. It is our hope that this list will serve as a point of reference for future studies and checklists; our goal will be met when arrangements adopted herein are proved incorrect through future research.

Synonyms, homonyms, and subspecies are not listed. A complete catalog of North American butterflies including all synonyms and subspecies is in preparation by Jonathan P. Pelham (Seattle, Washington). Citations to original descriptions have been provided for some recently described taxa (see References); these are indicated in the list by an asterisk. We have tried to be as precise as possible in our listing of taxon authors and dates of publication, and thank Gerardo Lamas (Lima, Peru) and J. Pelham for a great amount of help with these issues. All changes (from other checklists) in authorship or dates of publication made herein will be fully explained in Pelham's upcoming catalog. We note here, however, that our spelling of Leconte's name follows Rehn (1954).

This being a list of species, a few words on the species concepts held by the authors seem warranted. There are currently no fewer than 25 species concepts being used by taxonomists in various fields (Mayden 1997, Brower 1999, Sperling in press), and no fewer than five species concepts are currently applied in morphological and molecular studies of Lepidoptera taxonomy. Some of the more popular species concepts held by taxonomists today don't even recognize subspecies. While subspecies are frequently named to describe geographical variants, the criteria for recognizing such variants as "subspecies" are subjective and vary greatly from author to author (even when those authors claim to apply the same species concept).

The authors of this list apply two different species concepts when making taxonomic decisions. The senior author (PAO) holds views most consistent with Mayr's Biological Species Concept (Mayr 1957a,b). The most current version of this concept defines species as "groups of

interbreeding natural populations that are reproductively isolated from other such groups” (see Wheeler & Meier 2000). Earlier versions of Mayr’s concept required users to infer whether or not allopatric populations are capable of interbreeding, but later versions of the concept have abandoned this notion (since it is often impossible to make this inference anything other than a guess). The production of fertile offspring is an indication of a biological species.

The views of the junior author (ADW) are most consistent with the Phylogenetic Species Concept (Nixon & Wheeler 1990) when making taxonomic decisions. A phylogenetic species is defined as the “smallest aggregation of (sexual) populations or (asexual) lineages diagnosable by a unique combination of character states in comparable individuals (semaphoronts).” Under this concept, species are identified by unique character states, which show no intermediacy or signs of introgression with related organisms. For a debate between proponents of these and other species concepts, see Wheeler & Meier (2000).

Even with different species concepts, the authors of this list agreed on over 90% of the taxonomic decisions that have been incorporated herein, even before initial discussion. Subsequent discussion and re-evaluation of available evidence has resolved almost all the discrepancies in taxonomic opinions between the authors (only a few cases remain where the authors don’t entirely agree). Therefore, the majority of the species listed herein can be considered biological and phylogenetic species, in cases where sufficient information is known (and most likely qualify as species under other species concepts). Annotations after species entries will hopefully explain differences between this listing and previous checklists, and enable users with alternate species concepts to extract necessary information from this list.

SUPERFAMILY HESPERIOIDEA, FAMILY HESPERIIDAE

[Our arrangement of this family follows Evans (1951, 1952, 1953, 1955) and Voss (1952), as modified by Ackery et al. (1999) and Warren (2000, 2002). The higher-level classification of the family is currently being studied by ADW.]

Subfamily Pyrrhopyginae

Apyrrothrix Lindsey, 1921

1. *araxes* (Hewitson, 1867)

[Burns & Janzen (2001) treat *arizonae* (Godman & Salvin, 1893) as a full species based on genitalic differences observed in eleven male specimens, compared from Arizona and the Mexico City area. However, populations from areas in between (Sinaloa and Nayarit) show intermediacy in all of the genitalic differences cited by Burns and Janzen, and several additional geographic variants are known (ADW, unpublished). See Mielke (2002) for generic change from *Pyrrhopuge* Hübner, [1819].]

Subfamily Pyrginae

Phocides Hübner, [1819]

2. *pigmalion* (Cramer, 1779)

2.1 *belus* Godman & Salvin, 1893

Observed and photographed at Bentsen-Rio Grande State Park, Hidalgo Co. Texas on April 13, 2003 by David J. Hanson. Photo identified by Andrew D. Warren (Hanson, Knudson, & Bordelon, 2003).

3. *polybius* (Fabricius, 1793)

[*Papilio palemon* Cramer, 1777, being a primary homonym of *Papilio palaemon* Pallas, 1771, is not an available name (ICZN 1999).]

Proteides Hübner, [1819]

4. *mercurius* (Fabricius, 1787)

Epargyreus Hübner, [1819]

5. *clarus* (Cramer, 1775)

6. *zestos* (Geyer, 1832)

7. *exadeus* (Cramer, 1779)

8. species

[Arizona records of “*exadeus*” almost certainly refer to *E. windi* H.A. Freeman, 1969* (a resident of central Sonora, Mexico), but specimens have not been examined; *exadeus* is unknown in western Mexico north of Nayarit (ADW, unpublished). *Epargyreus aspina* Evans, 1952* could possibly also stray into our area; dissection of all exotic *Epargyreus* specimens in our area is needed to confirm their specific identity.]

Polygonus Hübner, [1825]

9. *leo* (Gmelin, [1790])

10. *savigny* (Latreille, [1824])

[Mielke & Casagrande (2002) treat *manueli* Bell & W.P. Comstock, 1948 as a junior

synonym of *savigny*; also see Mielke (2004).]

Chioides Lindsey, 1921

11. *albofasciatus* (Hewitson, 1867)

[Raised to species level by Austin and Warren (2002); previously considered to be a subspecies of *C. catillus* (Cramer, 1779).]

12. *zilpa* (Butler, 1872)

Aguna R.C. Williams, 1927

13. *asander* (Hewitson, 1867)

14. *claxon* Evans, 1952*

15. *metophis* (Latreille, [1824])

Typhedanus Butler, 1870

16. *undulatus* (Hewitson, 1867)

Polythrix E.Y. Watson, 1893

17. *octomaculata* (Sepp, [1844])

18. *mexicanus* H.A. Freeman, 1969*

Zestusa Lindsey, 1925

19. *dorus* (W.H. Edwards, 1882)

Codatractus Lindsey, 1921

20. *alcaeus* (Hewitson, 1867)

21. *arizonensis* (Skinner, 1905)

22. *valeriana* (Plötz, 1881)

[This species was placed in *Codatractus* by Burns (1996) as *mysie* (Dyar, 1904), but examination of the type of *valeriana* (Plötz) showed it to be the senior objective synonym (Mielke & Warren, 2004).]

Urbanus Hübner, [1807]

23. *proteus* (Linnaeus, 1758)

24. *belli* (Hayward, 1935)

[Specimen reported from south Texas by Warren (1997).]

25. *pronus* Evans, 1952*

26. *esmeraldus* (Butler, 1877)

27. *dorantes* (Stoll, 1790)

28. *teleus* (Hübner, 1821)

29. *tanna* Evans, 1952*

30. *simplicius* (Stoll, 1790)

31. *procne* (Plötz, 1881)

32. *doryssus* (Swainson, 1831)

Astraptes Hübner, [1819]

33. *fulgerator* (Walch, 1775)

34. *egregius* (Butler, 1870)

35. *alardus* (Stoll, 1790)

36. *alector* (C. Felder & R. Felder, 1867)

[*Astraptes gilberti* H.A. Freeman, 1969* is treated as a synonym of *A. alector hopfferi* (Plötz, 1881) by Scott (1986) and Warren (2000, 2002).]

37. *anaphus* (Cramer, 1777)

Autochton Hübner, 1823

38. *cellus* (Boisduval & Leconte, [1837])
 39. *pseudocellus* (Coolidge & Clemence, [1910])
[Apparently extirpated from the U.S portion of its range (Bailowitz & Brock 1991).]
 40. *cincta* (Plötz, 1882)

Achalarus Scudder, 1872

41. *lyciades* (Geyer, 1832)
 42. *casica* (Herrick-Schäffer, 1869)
 43. *albociliatus* (Mabille, 1877)
 44. *toxeus* (Plötz, 1882)

Thessia Steinhauser, 1989*

45. *jalapus* (Plötz, 1881)

Thorybes Scudder, 1871

46. *drusius* (W.H. Edwards, [1884])
 47. *pylades* (Scudder, 1870)
 48. *bathyllus* (J.E. Smith, 1797)
 49. *confusis* Bell, 1922
 50. *diversus* Bell, 1927
 51. *mexicana* (Herrick-Schäffer, 1869)

Cabares Godman & Salvin, 1894

52. *potrillo* (Lucas, 1857)

Celaenorrhinus Hübner, [1819]

53. *fritzgaertneri* (Bailey, 1880)
 54. *stallingsi* H.A. Freeman, 1946*

Spathilepia Butler, 1870

55. *clonius* (Cramer, 1775)

Cogia Butler, 1870

56. *hippalus* (W.H. Edwards, 1882)
 57. *outis* (Skinner, 1894)
 58. *caicus* (Herrick-Schäffer, 1869)
 59. *calchas* (Herrick-Schäffer, 1869)

Arteurotia Butler & H. Druce, 1872

60. *tractipennis* Butler & H. Druce, 1872

Nisoniades Hübner, [1819]

61. *rubescens* (Möschler, 1877)

Pellicia Herrich-Schäffer, 1870

62. *arina* Evans, 1953*

63. *dimidiata* Herrich-Schäffer, 1870

Noctuana Bell, 1937

64. *stator* (Godman, 1899)

[Documented from Hidalgo Co., Texas, by Bob Stewart (Anonymous 2000).]

Windia H.A. Freeman, 1969*

65. *windi* H.A. Freeman, 1969*

**[Specimen collected in Guadalupe Canyon, Cochise Co., Arizona, by Kilian Roever.
 Determined by J. Burns.]**

Bolla Mabille, 1903

66. *brennus* (Godman & Salvin, 1896)
 67. *clytius* (Godman & Salvin, 1897)
- Staphylus* Godman & Salvin, 1896
 68. *ceos* (W.H. Edwards, 1882)
 69. *mazans* (Reakirt, [1867])
 70. *hayhurstii* (W.H. Edwards, 1870)
- Gorgythion* Godman & Salvin, 1896
 71. *begga* (Prittwitz, 1868)
[Listed as *G. vox* Evans, 1953* by Neck (1996), although the U.S. specimens have been confirmed as *begga*. Dissection is required for certain identification. Also see Stanford (2002).]
- Sostrata* Godman & Salvin, 1895
 72. *nordica* Evans, 1953*
[Treatment as a full species follows Warren (2000, 2002); *S. bifasciata* (Ménétriés, 1829) is a South American species. The relationship between these two taxa is currently being studied in detail by G.T. Austin and ADW.]
- Carrhenes* Godman & Salvin, 1895
 73. *canescens* (R. Felder, 1869)
- Xenophanes* Godman & Salvin, 1895
 74. *tryxus* (Stoll, 1780)
[The species name is often misspelled (e.g. Neck 1996).]
- Antigonus* Hübner, [1819]
 75. *emorsa* (R. Felder, 1869)
 75.1 *erosus* (Hübner, 1812)
[*Antigonus erosus* was first reported from the United States by Knudson et al. 2004.).]
- Mylon* Godman & Salvin, 1885
 75.2 *pelopidas* (Fabricius, 1793)
[First reported from the United States by Davis et al. 2005.]
- Systasea* W.H. Edwards, 1877
 76. *pulverulenta* (R. Felder, 1869)
 77. *zampa* (W.H. Edwards, 1876)
- Achlyodes* Hübner, [1819]
 77.1 *pallida* (R. Felder, 1869)
[Photographed in the lower Rio Grande Valley in October, 2003 and reported by Warren et al., 2003.]
- Eantis* Boisduval, 1836
[Our treatment of *Eantis* is based on Warren (1996); these species were previously placed in *Achlyodes* Hübner, [1819]. Also see Brown & Heineman (1972).]
 78. *tamenund* (W.H. Edwards, 1871)
[*Eantis thraso* (Hübner, [1807]) is a Central and South American species (Warren 1996).]
- Grais* Godman & Salvin, 1894
 79. *stigmaticus* (Mabille, 1883)

Timochares Godman & Salvin, 1896

80. *ruptifasciata* (Plötz, 1884)

Anastrus Hübner, [1824]

80.1 *sempiternus* (Butler & H. Druce, 1872)

[***Anastrus sempiternus* (Butler and Druce) was first collected in Starr Co., Texas by Charles Bordelon (Knudson, 2002).**]

Chiomara Godman & Salvin, 1899

81. *georgina* (Reakirt, 1868)

[***Chiomara asychis* (Stoll, 1780) is a South American species that does not occur in Costa Rica (Janzen et al. 1998) or Mexico (Warren 2000, 2002); however, its relationship to *georgina* is currently being studied in detail by G. Austin and ADW.**]

82. *mithrax* (Möschler, 1879)

Gesta Evans, 1953

83. *invisus* (Butler & H. Druce, 1872)

[**Austin and Warren (2002) treat *G. invisus* and *G. gesta* (Herrich-Schäffer, 1863), a West Indian and South American taxon, as separate species.**]

Ephyriades Hübner, [1819]

84. *brunnea* (Herrich-Schäffer, 1865)

Erynnis Schrank, 1801

85. *icelus* (Scudder & Burgess, 1870)

86. *brizo* (Boisduval & Leconte, [1837])

[**Research is needed to determine if one or more subspecies should be elevated to the species-level.]**

87. *juvenalis* (Fabricius, 1793)

[**Research is needed to determine if *clitus* (W.H. Edwards, 1882) should be elevated to the species-level.]**

88. *telemachus* Burns, 1960

89. *propertius* (Scudder & Burgess, 1870)

90. *meridianus* Bell, 1927

91. *scudderii* (Skinner, 1914)

92. *horatius* (Scudder & Burgess, 1870)

93. *tristis* (Boisduval, 1852)

94. *martialis* (Scudder, [1870])

95. *pacuvius* (Lintner, 1878)

96. *zarucco* (Lucas, 1857)

97. *funeralis* (Scudder & Burgess, 1870)

98. *baptisiae* (Forbes, 1936)

99. *lucilius* (Scudder & Burgess, 1870)

100. *afranius* (Lintner, 1878)

101. *persius* (Scudder, 1863)

[**Research is needed to better define relationships among and within the above three taxa; see Burns (1964).]**

Pyrgus Hübner, [1819]

102. *centaureae* (Rambur, [1842])

[**Tentatively includes *wyandot* (W.H. Edwards, 1863), which was treated as a**

species-level taxon by Shapiro (1974) and Gochfield & Burger (1997) (based on genitalic differences between it and *freija* (Warren, 1924) cited by Forbes (1960); but see Lindsey (1928)). Its relationship to other North American and Eurasian taxa associated with *centaureae* remains to be studied in detail. Also see Warren (1926, 1935, 1951), de Jong (1972), Devyatkin (1990) and Tuzov et al. (1997).]

- 103. *ruralis* (Boisduval, 1852)
- 104. *xanthus* W.H. Edwards, 1878
- 105. *scriptura* (Boisduval, 1852)
- 106. *communis* (Grote, 1872)
- 107. *albescens* Plötz, 1884

[Treatment as a full species follows Burns (2001).]

- 108. *philetas* W.H. Edwards, 1881
- 109. *oileus* (Linnaeus, 1767)

[The above four taxa were placed in *Syrichtus* Boisduval, [1834] by Durden (1982). This is incorrect because *Syrichtus* is a synonym of *Pyrgus* (see Cowan 1970). There is no published evidence indicating that these four species belong in a Palaearctic genus other than *Pyrgus* (see de Jong 1972, 1978); however, further study is needed.]

Heliopyrgus Herrera, 1957

[Treatment of *Heliopyrgus* is based on Austin & Warren (2001); also see Mielke & Casagrande (2002).]

- 110. *domicella* (Erichson, [1849])

Heliopetes Billberg, 1820

- 111. *ericetorum* (Boisduval, 1852)
- 112. *macaira* (Reakirt, [1867])
- 113. *laviana* (Hewitson, 1868)
- 114. *arsalte* (Linnaeus, 1758)
- 114.1 *sublinea* (Schaus, 1902)

[First report for the United States by Basham et al., 2005]

Celotes Godman & Salvin, 1899

- 115. *nessus* (W.H. Edwards, 1877)
- 116. *limpia* Burns, 1974*

Pholisora Scudder, 1872

- 117. *catullus* (Fabricius, 1793)
- 118. *mejicanus* (Reakirt, [1867])

Hesperopsis Dyar, 1905

- 119. *libya* (Scudder, 1878)
- 120. *alpheus* (W.H. Edwards, 1876)
- 121. *gracielae* (MacNeill, 1970)*

Subfamily Heteropterinae

Carterocephalus Lederer, 1852

122. *palaemon* (Pallas, 1771)

[The relationships between New and Old World taxa of *palaemon* remain to be studied in detail.]

Piruna Evans, 1955*

123. *pirus* (W.H. Edwards, 1878)

124. *haferniki* H.A. Freeman, 1970*

125. *polingii* (Barnes, 1900)

126. *aea* (Dyar, 1912)

[Previously known as *P. cingo* Evans, 1955*, which was treated as a synonym of *aea* by Warren (2000, 2002).]

127. *penaea* (Dyar, 1918)

[Previously misidentified as *P. microsticta* (Godman, 1900) (see McGuire & Rickard 1976), a southwestern Mexican species (Freeman & Warren, in preparation).]

Subfamily Hesperiinae

[Includes genera formerly placed in the Megathyminae, following Ackery et al. (1999).]

Synapte Mabille, 1904

128. *pecta* Evans, 1955*

[Warren (2000, 2002) treated *S. pecta* and *S. malitiosa* (Herrich-Schäffer, 1865) as separate species.]

129. species

[We are uncertain if it is *Synapte syraces* (Godman, 1901), or the similar *S. shiva* Evans, 1955* (or both?) that is known from Arizona (Bailowitz & Brock 1991), since we have not examined specimens; both species occur in Sonora, Mexico. *Synapte shiva* was described as a subspecies of *syraces*, but was treated as a full species by Warren (2000, 2002).]

130. *salenus* (Mabille, 1883)

Corticea Evans, 1955*

131. *corticea* (Plötz, 1882)

Vidius Evans, 1955*

132. *perigenes* (Godman, 1900)

Monca Evans, 1955*

133. *crispinus* (Plötz, 1882)

[Mielke and Casagrande (2002) treat *tyrtaeus* (Plötz, 1882) as a junior synonym of *crispinus*. *Monca telata* (Herrich-Schäffer, 1869) is a South and Central American species (Bell 1941) which flies sympatrically with *crispinus* (= *tyrtaeus*) in Guatemala (Austin et al. 1996).]

Nastra Evans, 1955*

134. *Iherminier* (Latreille, [1824])

135. *neamathla* (Skinner & R.C. Williams, 1923)

136. *julia* (H.A. Freeman, 1945)

Cymaenes Scudder, 1872

137. *tripunctus* (Herrich-Schäffer, 1865)

138. *trebius* (Mabille, 1891)

[*Cymaenes trebius* was treated as a species distinct from *C. odilia* (Burmeister, 1878) by Warren (2000, 2002); however, relationships among taxa placed with *odilia* by Evans (1955) are under study by G. Austin and ADW.]

Lerema Scudder, 1872

139. *accius* (J.E. Smith, 1797)

140. *liris* Evans, 1955*

[The relationship between *L. liris* and *L. ancillaris* (Butler, 1877) where they meet in Central America needs further study, as noted by Warren (2000).]

Vettius Godman, 1901

141. *fantasos* (Cramer, 1780)

Perichares Scudder, 1872

142. *philetis* (Gmelin, [1790])

Rhinthon Godman, 1900

143. *osca* (Plötz, 1882)

[Treatment as a distinct species from *cubana* (Herrich-Schäffer, 1865) follows McGuire & Rickard (1976), Ferris (1989c) and Warren (2000, 2002), but see Mielke & Casagrande (2002).]

Decinea Evans, 1955*

144. *percosius* (Godman, 1900)

Conga Evans, 1955*

145. *chydaea* (Butler, 1877)

Ancyloxypha C. Felder, 1862

146. *numitor* (Fabricius, 1793)

147. *arene* (W.H. Edwards, 1871)

Oarisma Scudder, 1872

148. *garita* (Reakirt, 1866)

149. *powesheik* (Parker, 1870)

[Corrected to the original orthography even though name was misspelled.]

150. *edwardsii* (Barnes, 1897)

Copaeodes Speyer, 1877

151. *aurantiaca* (Hewitson, 1868)

152. *minima* (W.H. Edwards, 1870)

Adopaeoides Godman, 1900

153. *prittwitti* (Plötz, 1884)

Thymelicus Hübner, [1819]

154. *lineola* (Ochsenheimer, 1808)

Hylephila Billberg, 1820

155. *phyleus* (Drury, 1773)

Pseudocopaeodes Skinner & R.C. Williams, 1923

156. *eunus* (W.H. Edwards, 1881)

Stinga Evans, 1955*

157. *morrisoni* (W.H. Edwards, 1878)

Hesperia Fabricius, 1793

158. *uncas* W.H. Edwards, 1863

- 159. *juba* (Scudder, 1874)
- 160. *comma* (Linnaeus, 1758)
- 161. *assiniboia* (Lyman, 1892)

[Species-level status follows Layberry et al. (1998), but further elaboration on the relationship between this taxon and the next is needed.]

- 162. *colorado* (Scudder, 1874)

[*Hesperia colorado* (Scudder, 1874) has priority over *H. harpalus* (W.H. Edwards, 1881) and *H. idaho* (W.H. Edwards, 1883). The *Hesperia comma* complex was discussed by MacNeill (1964, 1975) and by Scott (1975, 1998). Relationships of taxa placed with *assiniboia*, *comma*, and *colorado* are currently being studied by several researchers.]

- 163. *woodgatei* (R.C. Williams, 1914)
- 164. *ottoe* W.H. Edwards, 1866
- 165. *leonardus* Harris, 1862
- 166. *pahaska* (Leussler, 1938)
- 167. *columbia* (Scudder, 1872)
- 168. *metea* Scudder, 1864
- 169. *viridis* (W.H. Edwards, 1883)
- 170. *attalus* (W.H. Edwards, 1871)
- 171. *meskei* (W.H. Edwards, 1877)
- 172. *dacotae* (Skinner, 1911)
- 173. *lindseyi* (Holland, 1930)
- 174. *sassacus* Harris, 1862
- 175. *miriamae* MacNeill, 1959
- 176. *nevada* (Scudder, 1874)

Atalopedes Scudder, 1872

- 177. *campestris* (Boisduval, 1852)

Polites Scudder, 1872

[See Burns (1994a) for synonymy of *Yvretta* Hemming, 1935 under *Polites*.]

- 178. *rhesus* (W.H. Edwards, 1878)
- 179. *carus* (W.H. Edwards, 1883)
- 180. *peckius* (W. Kirby, 1837)
- 181. *sabuleti* (Boisduval, 1852)
- 182. *draco* (W.H. Edwards, 1871)
- 183. *mardon* (W.H. Edwards, 1881)
- 184. *themistocles* (Latreille, [1824])
- 185. *baracoa* (Lucas, 1857)
- 186. *origenes* (Fabricius, 1793)
- 187. *mystic* (W.H. Edwards, 1863)
- 188. *sonora* (Scudder, 1872)
- 189. *vibex* (Geyer, 1832)

Wallengrenia Berg, 1897

- 190. *otho* (J.E. Smith, 1797)

[Tentatively includes *clavus* (Erichson, [1849]) and *curassavica* (Snellen, 1887); see

Burns (1994b). The complex is being studied by J.Y. Miller.]

191. *egeremet* (Scudder, 1864)

Pompeius Evans, 1955*

192. *verna* (W.H. Edwards, 1862)

Atrytone Scudder, 1872

193. *arogos* (Boisduval & Leconte, [1837])

Anatrytone Dyar, 1905

[Generic combination follows Burns (1994b).]

194. *logan* (W.H. Edwards, 1863)

195. *mazai* (H.A. Freeman, 1969)*

Problema Skinner & R.C. Williams, 1924

196. *byssus* (W.H. Edwards, 1880)

197. *bulenta* (Boisduval & Leconte, [1837])

Ochlodes Scudder, 1872

198. *sylvanoides* (Boisduval, 1852)

199. *agricola* (Boisduval, 1852)

200. *yuma* (W.H. Edwards, 1873)

[Our treatment of *Ochlodes* species follows Chiba and Tsukiyama (1996).]

Poanes Scudder, 1872

201. *hobomok* (Harris, 1862)

202. *zabulon* (Boisduval & Leconte, [1837])

203. *taxiles* (W.H. Edwards, 1881)

204. *melane* (W.H. Edwards, 1869)

205. *massasoit* (Scudder, 1864)

206. *viator* (W.H. Edwards, 1865)

207. *yehl* (Skinner, 1893)

208. *aaroni* (Skinner, 1890)

Paratrytone Godman, 1900

[See Burns (1992) for generic combination.]

209. *snowi* (W.H. Edwards, 1877)

Quasimellana Burns, 1994*

210. *eulogius* (Plötz, 1882)

Euphyes Scudder, 1872

211. *pilatka* (W.H. Edwards, 1867)

212. *conspicua* (W.H. Edwards, 1863)

213. *berryi* (Bell, 1941)

214. *dion* (W.H. Edwards, 1879)

[Includes *E. alabamae* (Lindsey, 1923) as a subspecies, and *E. macguirei* H.A. Freeman, 1975* as a synonym, following Shuey (1989); also see Shuey (1993).]

215. *bayensis* Shuey, 1989*

216. *dukesii* (Lindsey, 1923)

217. *bimacula* (Grote & Robinson, 1867)

218. *arpa* (Boisduval & Leconte, [1837])

219. *vestris* (Boisduval, 1852)

Asbolis Mabille, 1904

220. *capucinus* (Lucas, 1857)

Atrytonopsis Godman, 1900

221. *hianna* (Scudder, 1868)

222. *loammi* (Whitney, 1876)

223. *deva* (W.H. Edwards, 1877)

224. *lunus* (W.H. Edwards, 1884)

225. *vierecki* (Skinner, 1902)

226. *pittacus* (W.H. Edwards, 1882)

227. *python* (W.H. Edwards, 1882)

228. *cestus* (W.H. Edwards, 1884)

229. *edwardsi* Barnes & McDunnough, 1916

Amblyscirtes Scudder, 1872

230. *exoteria* (Herrich-Schäffer, 1869)

231. *cassus* W.H. Edwards, 1883

232. *aenus* W.H. Edwards, 1878

[Includes *erna* H.A. Freeman, 1943, following Scott (1977).]

233. *linda* H.A. Freeman, 1943

234. *oslari* (Skinner, 1899)

235. *elissa* Godman, 1900

236. *hegon* (Scudder, 1864)

237. *texanae* Bell, 1927

238. *carolina* (Skinner, 1892)

239. *reversa* W.M. Jones, 1926

240. *aesculapius* (Fabricius, 1793)

241. *nereus* (W.H. Edwards, 1876)

242. *nysa* W.H. Edwards, 1877

243. *eos* (W.H. Edwards, 1871)

244. *vialis* (W.H. Edwards, 1862)

245. *alternata* (Grote & Robinson, 1867)

246. *celia* Skinner, 1895

247. *belli* H.A. Freeman, 1941

248. *tolteca* Scudder, 1872

[Includes *prenda* Evans, 1955*, following Freeman (1993).]

249. *phylace* (W.H. Edwards, 1878)

250. *fimbriata* (Plötz, 1882)

251. *simius* W.H. Edwards, 1881

[While Burns (1990) removed this species from *Amblyscirtes* and left it “*incertae sedis*,” we list *simius* here until a generic placement is proposed.]

Lerodea Scudder, 1872

252. *eufala* (W.H. Edwards, 1869)

253. *arabus* (W.H. Edwards, 1882)

[Includes *dysaules* Godman, 1900 as a synonym (Warren 2000, 2002; Warren & Mielke 2005).]

Oligoria Scudder, 1872

254. *maculata* (W.H. Edwards, 1865)

- Calpodes* Hübner, [1819]
 255. *ethlius* (Stoll, 1782)
- Panoquina* Hemming, 1934*
 256. *panoquin* (Scudder, 1864)
 257. *panoquinoides* (Skinner, 1891)
 258. *errans* (Skinner, 1892)
 259. *ocola* (W.H. Edwards, 1863)
 260. *lucas* (Fabricius, 1793)
 [*Panoquina sylvicola* (Herrick-Schäffer, 1865) was treated as a synonym of *lucas* by Robbins et al. (1996); also see Mielke & Casagrande (2002).]
 261. *hecebolum* (Scudder, 1872)
 262. *evansi* (H.A. Freeman, 1946)*
- Nyctelius* Hayward, 1948
 263. *nyctelius* (Latreille, [1824])
- Thespieus* Godman, 1900
 264. *macareus* (Herrick-Schäffer, 1869)
- Agathymus* H.A. Freeman, 1959*
 265. *neumoegeni* (W.H. Edwards, 1882)
 [Includes *chisosensis* (H.A. Freeman, 1952), following Roever (1975).]
 266. *polingi* (Skinner, 1905)
 267. *evansi* (H.A. Freeman, 1950)
 268. *aryxna* (Dyar, 1905)
 269. *baueri* (D. Stallings & Turner, 1954)
 270. *gentryi* Roever, 1998*
 [Treatment of this taxon and *baueri* as subspecies of *aryxna* is unsupported by any compelling data (Cassie et al. 2001), and does not consider problems associated with the name *aryxna*; see dos Passos (1960) and Roever (1975).]
 271. *mariae* (Barnes & Benjamin, 1924)
 [Tentatively includes *gilberti* H.A. Freeman, 1964*, although the status of this taxon is uncertain (see Roever 1975).]
 272. *estelleae* (D. Stallings & Turner, 1958)
 [Includes *valverdiensis* H.A. Freeman, 1966, following Roever (1975).]
 273. *stephensi* (Skinner, 1912)
 274. *alliae* (D. Stallings & Turner, 1957)
- Megathymus* Scudder, 1872
 275. *yuccae* (Boisduval & Leconte, [1837])
 [Includes *coloradensis* C.V. Riley, 1877, following Roever (1975).]
 276. *streckeri* (Skinner, 1895)
 277. *cofaqui* (Strecker, 1876)
 278. *ursus* Poling, 1902
- Stallingsia* H.A. Freeman, 1959*
 279. *maculosus* (H.A. Freeman, 1955)

SUPERFAMILY PAPILIONOIDEA

FAMILY PAPILIONIDAE

[We have followed the generic arrangement proposed for this family by Miller (1987). We feel further study is needed on the generic relationships of swallowtails in order to subdivide them into smaller monophyletic groups with confidence.]

Subfamily Parnassiinae

Parnassius Latreille, 1804

280. *eversmanni* [Ménétriés], [1850]

281. *clodius* Ménétriés, 1857

282. *phoebus* (Fabricius, 1793)

283. *behrii* W.H. Edwards, 1870

284. *smintheus* Doubleday, [1847]

[Split of *P. phoebus* complex is discussed by Shepard & Manley (1998), Bird et al. (1995) and Layberry et al. (1998).]

Subfamily Papilioninae

Battus Scopoli, 1777

285. *philenor* (Linnaeus, 1771)

286. *polydamas* (Linnaeus, 1758)

Parides Hübner, [1819]

287. *eurimedes* (Stoll, 1782)

288. *alopius* (Godman & Salvin, 1890)

[Single specimen collected in Cochise Co., Arizona, by Neil Dankert; see Bailowitz & Brock (1991).]

Eurytides Hübner, [1821]

289. *marcellus* (Cramer, 1777)

290. *philolaus* (Boisduval, 1836)

Papilio Linnaeus, 1758

291. *machaon* Linnaeus, 1758

[Includes *bairdii* W.H. Edwards, 1866, and *oregonius* W.H. Edwards, 1876, following Sperling (1987, 1993a, 2003) and Sperling and Harrison (1994); but see Eitschberger (1993) and Pyle (2002).]

292. *brevicauda* Saunders, 1869

293. *joanae* Heitzman, 1973*

294. *polyxenes* Fabricius, 1775

295. *zelicaon* Lucas, 1852

[*Papilio nitra* W.H. Edwards, 1883 is considered by Fisher (1977) to be an eastern subspecies of *zelicaon* containing rare dark individuals, which may represent infrequent introgression with *polyxenes*. Lamas (2004) considers this to be a subspecies of *P. polyxenes* Fabricius despite a lack of published evidence of conspecificity.]

296. *indra* Reakirt, 1866

297. *glaucus* Linnaeus, 1758

297.1 *appalachensis* (Pavulaan & Wright, 2002)

[*Pterourus appalachiensis* Pavulaan & D. Wright, 2002* was described, and suggestive evidence for its treatment as a full species was provided by Scriber and Ording, 2005.]

298. *canadensis* Rothschild & Jordan, 1906

[Specific distinctness of *P. canadensis* is detailed by Hagen et al. (1991), and further supported by Sperling (1993b).]

299. *rutulus* Lucas, 1852

300. *eurymedon* Lucas, 1852

301. *multicaudata* W.F. Kirby, 1884

302. *pilumnus* Boisduval, 1836

303. *troilus* Linnaeus, 1758

304. *palamedes* Drury, 1773

305. *garamas* (Geyer, [1829])

[*Papilio garamas abderus* Hoppfer, 1856 was reported from South Texas by Bordelon & Knudson (2000).]

306. *victorinus* Doubleday, 1844

[Treated as a subspecies of *P. menatius* Hübner [1819] by Tyler et al. (1994: but see page 26) and Lamas (2004); however, we feel that further study of the situation is needed (see DeVries 1987). The species was first reported from the U.S. by Adams (1984).]

307. *thoas* Linnaeus, 1771

308. *cresphontes* Cramer, 1777

309. *astyalus* Godart, 1819

310. *ornythion* Boisduval, 1836

311. *aristodemus* Esper, 1794

312. *andraemon* (Hübner, [1823])

313. *androgeus* Cramer, 1775

314. *anchisiades* Esper, 1788

315. *rogeri* Boisduval, 1836

[*Papilio pharnaces* Doubleday, 1846 is treated as a subspecies of *P. rogeri* Boisduval, 1836 by Lamas (2004); the two taxa were also considered to be conspecific by Tyler et al. (1994).]

*[Miller (1987) does not recognize *Heraclides* Hübner [1819], *Priamides* Hübner [1819] or *Pterourus* Scopoli, 1777 as valid genera. Tyler et al. (1994) present a considerably different generic arrangement for our swallowtail taxa; also see Lamas (2004).]

FAMILY PIERIDAE

Subfamily Dismorphiinae

Enantia Hübner, [1819]

316. *albania* (H.W. Bates, 1864)

[Texas specimen of this species illustrated by Kendall (1974a); see Llorente (1984) for notes on *Enantia* taxonomy.]

Subfamily Pierinae

Catasticta Butler, 1870

317. *nimbice* (Boisduval, 1836)

Neophasia Behr, 1869

318. *menapia* (C. Felder & R. Felder, 1859)

319. *terlooii* Behr, 1869

[Spelling corrected to original orthography, following Llorente et al. (1997) and Lamas (2004).]

Appias Hübner, [1819]

320. *drusilla* (Cramer, 1777)

[This species is often placed in the genus *Glutophrissa* Butler, 1887, e.g. Hemming (1967) and Lamas (1981, 2004); but see Klots (1933), d'Almeida (1939a), Comstock (1943) and Brown & Heineman (1972).]

Leptophobia Butler, 1870

321. *aripa* (Boisduval, 1836)

[Collected in Santa Ana National Wildlife Refuge, Hidalgo Co., Texas, by T. Kral (20-X-1988). Specimens deposited in C.P. Gillette Museum, Colorado State University.]

Pontia Fabricius, 1807

322. *beckerii* (W.H. Edwards, 1871)

[Results of studies by Chew & Watt (2006) show that *P. beckerii* is sufficiently divergent from *P. chloridice* (Hübner, [1813]).]

323. *protodice* (Boisduval & Leconte, [1830])

324. *occidentalis* (Reakirt, 1866)

[Chew and Watts find sufficient mtDNA divergence with *P. callidice* (Hübner, [1800]) to suggest that it is not conspecific with *occidentalis*.]

324.1 *nelsoni* (Edwards, 1883)

[Chew & Watts (2006) have shown that the mtDNA sequence of this taxon shows that it is strongly diverged from *P. occidentalis* (Reakirt).]

325. *sisymbrii* (Boisduval, 1852)

Pieris Schrank, 1801

326. *rapae* (Linnaeus, 1758)

327. *oleracea* Harris, 1829

328. *marginalis* Scudder, 1861

[Relationship of *meckyae* Eitschberger, 1983* to *angelika* and *marginalis* needs clarification. Chew and Watt (2006) find sufficient mtDNA evidence to raise *macdunnoughii* (Remington) to the species level while leaving *angelika* as a synonym of *oleracea*.]

329. *angelika* Eitschberger, 1983*

[Study of type specimens is necessary; this may be a synonym of *P. pseudobryoniae* Barnes & McDunnough, 1917, which is an elevation of *P. napi* variety *frigida* form *pseudobryoniae* Verity, [1908], fide J. Pelham.]

*[*Pieris napi* Linnaeus, 1758, a Palaearctic taxon, does not occur in North America as per several authors, e.g. Chew & Watt, 2006. North American species exclusive of the distinct *Pieris virginensis* are treated as either three species by Chew & Watt

(2006), three species by Eitschberger (1981, 1983) or four species by Geiger and Shapiro (1992). This entire complex needs much more study before relationships can be understood with confidence. Also see Bowden (1972), Shapiro (1984), and Kudrna & Geiger (1985).]

330. *virginiensis* (W.H. Edwards, 1870)

*[See Robbins & Henson (1986) and Chew & Watt (2006) for use of *Pieris* over *Artogeia* Verity, 1947.]

Ascia Scopoli, 1777

331. *monuste* (Linnaeus, 1764)

Ganya Billberg, 1820

332. *josephina* (Godart, 1819)

333. *howarthi* (Dixey, 1915)

[Bailowitz (1988) describes the distribution and biology of this species in contrast with *G. josephina*; also see Beutelspacher (1986) and Llorente et al. (1997).]

Melete Swainson, [1831]

333.1. *lycimnia* (Cramer, 1777)

[*Melete lycimnia isandra* (Boisduval), 1836—One individual was collected in November 2004 in Mission, Hidalgo County, Texas. It is resident as far north as Ciudad Victoria in Tamaulipas state, Mexico (Dauphin et al. 2005).]

Itaballia Kaye, 1904

333.2. *demophile* (Linnaeus, 1763)

[*Itaballia demophile centralis* Joicey and Talbot, 1928—One individual was photographed in Mission, Hidalgo County, Texas during December, 2004. The butterfly is native to Central and South America. It is resident only as far north as San Luis Potosi state in Mexico (Dauphin et al. 2005).]

Pieriballia Klots, 1933

333.3. *viardi* (Boisduval, 1836)

[Bordelon & Knudson (2006) document a female that was photographed by Booker, Gustafson, Dauphin, Stangeland, Davis, and others at Bentsen State Park/World Birding Center in Hidalgo County, Texas on December 6, 2005.]

Euchloe Hübner, [1819]

334. *ausonides* (Lucas, 1852)

[Includes *ogilvia* Back, [1991]*, although this was described as a full species; see Layberry et al. (1998).]

335. *naina* Kozhanchikov, 1923

[Reported for North America by Kondla and Pelham (1995). Specific identity is described by Dubatolov and Kosterin (1994).]

336. *guaymasensis* Opler, 1987*

[Specimen collected in Bisbee, Cochise Co., Arizona, by Sandy Upson. Identity verified by R. Bailowitz, PAO and ADW.]

337. *olympia* (W.H. Edwards, 1871)

338. *creusa* (Doubleday, [1847])

339. *hyantis* (W.H. Edwards, 1871)

340. *lotta* Beutenmüller, 1898

[Above two species separated on basis of research by Opler (1965, 1967a,b, 1970,

1971, 1974) as well as unpublished data and communications. Detailed investigation has been initiated by PAO.]

Anthocharis Boisduval, Rambur, [Duméril] & Graslin, [1833]

341. *cethura* C. Felder & R. Felder, 1865

[Includes *pima* W.H. Edwards, 1888. These two form a gradual cline between California, Nevada, and Arizona, as discussed by Emmel et al. (1998b: p. 132).]

342. *sara* Lucas, 1852

343. *julia* W.H. Edwards, 1872

344. *stella* W.H. Edwards, 1879

345. *thoosa* (Scudder, 1878)

[The *sara* species group is split, in part, on basis of research by Geiger and Shapiro (1986); however, the overall distributions of these taxa remain somewhat unclear and the above arrangement is tentative. The complex is currently under study by PAO.]

346. *lanceolata* Lucas, 1852

347. *midea* (Hübner, [1809])

*[*Paramidea* Kusnezov, 1929 was described as a monotypic genus for the Asian *scolymus* (Butler, 1866). It was applied to *lanceolata* and *midea* by Ferris (1989c) without justification. It has been treated by several authors as a subgenus of *Anthocharis*, and is herein excluded as relating to any North American species.]

Subfamily Coliadinae

Colias Fabricius, 1807

348. *philodice* Godart, 1819

349. *eurytheme* Boisduval, 1852

350. *occidentalis* Scudder, 1862

351. *christina* W.H. Edwards, 1863

[This complex includes (among others) *krauthii* Klots, 1935, *kluanensis* Ferris, 1981*, *pseudochristina* Ferris, 1989*, *astraea* (W.H. Edwards, 1872), and several enigmatic eastern Oregon populations (see Ferris 1993). Layberry et al. (1998) treat *occidentalis* and *christina* (which includes all populations that are variably UV reflective and containing some orange individuals) as separate species, due to Ferris' statement that the variable Oregon populations were highly localized. However, these Oregon populations have proven to be very widespread over a large geographic area, and show a mosaic of variable segregates with different average character combinations. The complex is currently under study by Paul Hammond and David McCorkle.]

352. *alexandra* W.H. Edwards, 1863

353. *harfordii* Hy. Edwards, 1877

[Status of *C. harfordii* is detailed by Burns (1975) and Ferris (1988, 1993).]

354. *meadii* W.H. Edwards, 1871

355. *johanseni* Troubridge & Philip, 1990*

356. *hecla* Lefèvre, 1836

357. *canadensis* Ferris, 1982

358. *tyche* Böber, 1812

[Includes *boothii* Curtis, 1835 and *thula* Hovanitz, 1955, following Lafontaine & Wood (1997) and Layberry et al. (1998).]

359. *nastes* (Boisduval, 1832)

360. *scudderii* Reakirt, 1865

361. *gigantea* Strecker, 1900

362. *pelidne* Boisduval & Leconte, [1830]

363. *interior* Scudder, 1862

364. *palaeno* (Linnaeus, 1761)

[North American populations of this species have been referred to as species *chippewa* W.H. Edwards, 1872 (see Tuzov et al. 1997, Verhulst 2000, and Guppy & Shepard 2001), restricting *palaeno* to the Palaearctic region. Opinions vary, however (see Gorbunov 2001) and until a revision of the group is presented we retain *palaeno* for our populations.]

365. *behrii* W.H. Edwards, 1866

Zerene Hübner, [1819]

366. *cesonia* (Stoll, 1790)

367. *eurydice* (Boisduval, 1855)

Anteos Hübner, [1819]

368. *maerula* (Fabricius, 1775)

369. *clorinde* (Godart, [1824])

Phoebis Hübner, [1819]

370. *sennae* (Linnaeus, 1758)

371. *argante* (Fabricius, 1775)

[Stray specimens are known from Kansas (Field 1940) and Texas (Neck 1996, Stanford 2002).]

372. *agarithe* (Boisduval, 1836)

373. *philea* (Linnaeus, 1763)

374. *neocypris* (Hübner, [1823])

Aphrissa Butler, 1873

375. *statira* (Cramer, 1777)

376. *orbis* (Poey, 1832)

*[Recognition of *Aphrissa* follows Brown (1931), d'Almeida (1939b), Smith et al. (1994), Llorente et al. (1997), and Lamas (2004); but see Klots (1933). Also see Brown (1929) and d'Almeida (1940). No recent information to support or refute the lumping of *Aphrissa* with *Phoebis* has been presented, but further research is warranted.]

Kricogonia Reakirt, 1863

377. *lyside* (Godart, 1819)

Eurema Hübner, [1819]

378. *daira* (Godart, 1819)

379. *boisduvaliana* (C. Felder & R. Felder, 1865)

[Treated as a subspecies of *E. arbela* Geyer, 1832, by Lamas (2004).]

380. *mexicana* (Boisduval, 1836)

381. *salome* (C. Felder & R. Felder, 1861)

382. *albula* (Cramer, 1775)

[First U.S. record reported by Chuah & Cushing (1995).]

Pyrisitia Butler, 1870

- 383. *messalina* (Fabricius, 1787)
- 384. *proterpia* (Fabricius, 1775)
- 385. *lisa* (Boisduval & Leconte, [1830])
- 386. *nise* (Cramer, 1775)
- 387. *dina* (Poey, 1832)

Abaeis Hübner, [1819]

- 388. *nicippe* (Cramer, 1779)

*[Although the above three genera are sometimes lumped under *Eurema* (e.g. Ferris 1989c), recent DNA analysis by Pollock et al. (1998) supports our arrangement.]

Nathalis Boisduval, 1836

- 389. *iole* Boisduval, 1836

FAMILY LYCAENIDAE

[Our arrangement of this family follows Ackery et al. (1999), which corroborates Eliot (1990), who unites the coppers, blues and hairstreaks into a single subfamily; also see Eliot (1973).]

Subfamily Miletinae

Feniseca Grote, 1869

- 390. *tarquinius* (Fabricius, 1793)

Subfamily Lycaeninae

Tribe Lycaenini (Coppers)

Lycaena Fabricius, 1807

[We place all of our coppers in the genus *Lycaena* following the lead of Klots (1936); also see Pratt et al. (1993) and Lamas (2004). We feel that further study is needed on the generic relationships of copper species worldwide (but see Miller & Brown 1979).]

- 391. *arota* (Boisduval, 1852)
- 392. *phlaeas* (Linnaeus, 1761)

[Relationships between eastern and western North American, Eurasian, and African populations of *phlaeas* need further study (see Larsen 1991).]

- 393. *cupreus* (W.H. Edwards, 1870)

[Tentatively includes *snowi* (W.H. Edwards, [1881]), although further elaboration on its status is needed.]

- 394. *xanthoides* (Boisduval, 1852)
- 395. *dione* (Scudder, 1868)
- 396. *editha* (Mead, 1878)
- 397. *gorgon* (Boisduval, 1852)
- 398. *hyllus* (Cramer, 1775)

[The name *hyllus* appears to be a *nomen dubium* (see ICZN, 1999), and the use of *thoe* (Guérin-Méneville, [1831]) in its place may be warranted; see Koçak (1983).]

- 399. *rubidus* (Behr, 1866)

- 400. *ferrisi* K. Johnson & Balogh, 1977*
- 401. *heteronea* Boisduval, 1852
- 402. *epixanthe* (Boisduval & Leconte, [1835])
- 403. *holloides* (Boisduval, 1852)
- 404. *dorcas* (W. Kirby, 1837)
- 405. *dospassosi* McDunnough, 1940

[Treated as a distinct species by Layberry et al. (1998) and Handfield (1999). Further study is needed to delineate this and related taxa (*dorcas* and *holloides*) with confidence; see Ferris (1977) and Scott (1978).]

- 406. *nivalis* (Boisduval, 1869)
- 407. *mariposa* (Reakirt, 1866)
- 408. *hermes* (W.H. Edwards, 1870)

Tribe Theclini (Hairstreaks)

- Hypaurotis* Scudder, 1876
- 409. *crysalus* (W.H. Edwards, 1873)

- Habrodais* Scudder, 1876
- 410. *grunus* (Boisduval, 1852)

Tribe Eumaeini (Hairstreaks)

[The order of genera within the Eumaeini follows Robbins (2004).]

- Eumaeus* Hübner, [1819]
- 411. *toxea* (Godart, [1824])

[The occurrence of this species in our area needs to be verified; see Kendall & McGuire (1984).]

- 412. *atala* (Poey, 1832)

- Atlides* Hübner, [1819]

- 413. *halesus* (Cramer, 1777)

- Callophrys* Billberg, 1820

- 414. *affinis* (W.H. Edwards, 1862)

[Tentatively includes *apama* (W.H. Edwards, 1882) and *homoperplexa* Barnes & Benjamin, 1923, in accord with the treatment of Scott (1986); however, further elaboration on the relationship between *affinis* and *homoperplexa* is very badly needed.]

- 415. *perplexa* Barnes & Benjamin, 1923

[*Callophrys perplexa* and *C. affinis* occur together in the Pacific Northwest while maintaining separate hosts, habitats, and larval characteristics, according to Pyle (2002); also see Hinchliff (1996).]

- 416. *dumetorum* (Boisduval, 1852)

[Previously known as *C. viridis* W.H. Edwards, 1862, a synonym with the same type locality (Emmel et al. (1998a). Warren (2005) considers this taxon to be conspecific with *C. sheridanii* (W.H. Edwards, but nomenclatural issues remain to be resolved.)]

- 417. *sheridanii* (W.H. Edwards, 1877)

[Includes *comstocki* Henne, 1940, and *lemberti* Tilden, 1963, partly on evidence presented by Austin (1998b). The complex is being studied by ADW, J. Pelham & R. Stanford.]

- 418. *hesseli* (Rawson & Ziegler, 1950)

419. *nelsoni* (Boisduval, 1869)

[Includes *C. rosneri* K. Johnson, 1976*; The status of *C. byrnei* K. Johnson, 1976* is uncertain (see Ferris 1991). Putative areas of contact between *nelsoni plicataria* K. Johnson, 1976* and *gryneus barryi* K. Johnson, 1976* need further study (see Hinchliff 1994).]

420. *thornei* (J.W. Brown, 1983)*

[Differences between this species and nearby juniper-feeding populations are detailed by Brown (1983, 1993).]

421. *muiri* (Hy. Edwards, 1881)

[Identity described by Tilden (1952), where it occurs parapatrically with *nelsoni*. Species status was further supported by Nice and Shapiro (2001).]

422. *gryneus* (Hübner, [1819])

[Includes all juniper-feeding taxa, including *barryi*, *loki* (Skinner, 1907), *siva* (W.H. Edwards, 1874), and *sweedneri* (Chermock, 1945), among others; see Ferris (1991).]

423. *spinetorum* (Hewitson, 1867)

[Includes *millerorum* Clench, 1981* as a synonym following Robbins (1990).]

424. *johsoni* (Skinner, 1904)

425. *xami* (Reakirt, [1867])

426. *mcfarlandi* P. Ehrlich & Clench, 1960

427. *augustinus* (Westwood, 1852)

[Guppy and Shepard (2001) treat *iroides* (Boisduval, 1852) as a distinct species but provide little evidence to support the split; also see Kondla (1999).]

428. *mossii* (Hy. Edwards, 1881)

429. *fotis* (Strecker, [1878])

430. *polios* (Cook & Watson, 1907)

431. *irus* (Godart, [1824])

432. *henrici* (Grote & Robinson, 1867)

[Includes *solatus* (Cook & Watson, 1909) as a subspecies, although this is sometimes treated as a full species (e.g. Durden 1982).]

433. *niphon* Hübner, [1819]

434. *eryphon* (Boisduval, 1852)

435. *lanoraieensis* (Sheppard, 1934)

*[We have followed Clench (1961) and Robbins (2004) in our generic arrangement of *Callophrys* (s.l.), and treat *Deciduphagus* Johnson, 1992*, *Loranthomitoura* Ballmer & Pratt, 1992*, *Xamia* Clench, 1961*, *Sandia* Clench & P. Ehrlich, 1960, *Mitoura* Scudder, 1872 and *Incisalia* Scudder, 1872 as subgenera; but see Ballmer & Pratt (1992) and Johnson (1992).]

Cyanophrys Clench, 1961*

436. *goodsoni* (Clench, 1946)

437. *herodotus* (Fabricius, 1793)

438. *miserabilis* (Clench, 1946)

Rekoa Kaye, 1904

439. *palegon* (Cramer, 1780)

440. *marius* (Lucas, 1857)

Arawacus Kaye, 1904

441. *jada* (Hewitson, 1867)

Satyrium Scudder, 1876

442. *favonius* (J.E. Smith, 1797)

443. *ilavia* (Beutenmüller, 1899)

444. *polingi* (Barnes & Benjamin, 1926)

445. *titus* (Fabricius, 1793)

446. *acadica* (W.H. Edwards, 1862)

447. *californica* (W.H. Edwards, 1862)

448. *sylvinus* (Boisduval, 1852)

[Includes *dryope* (W.H. Edwards, 1870).]

449. *caryaevorus* (McDunnough, 1942)

450. *edwardsii* (Grote and Robinson, 1867)

451. *calanus* (Hübner, [1809])

452. *kingi* (Klots & Clench, 1952)

453. *liparops* (Leconte, 1833)

454. *auretorum* (Boisduval, 1852)

455. *tetra* (W.H. Edwards, 1870)

456. *saepium* (Boisduval, 1852)

457. *behrii* (W.H. Edwards, 1870)

458. *fuliginosa* (W.H. Edwards, 1861)

458.1 *semiluna* Klots, 1930

[Warren (2005) discusses the status of both species and a paper is in preparation by Opler, Warren, and Austin. Several apparent relatives are found in temperate Asia (Tuzov et al. 2000, Weidenhoffer et al. 2004).]

*[Species previously placed in *Fixsenia* Tutt, 1907 and *Harkenclenus* dos Passos, 1970, are now placed in *Satyrium* by Robbins (2004).]

Phaeostrymon Clench, 1961*

459. *alcestis* (W.H. Edwards, 1871)

Ocaria Clench, 1970

460. *ocrisia* (Hewitson, 1868)

Chlorostrymon Clench, 1961*

461. *simaethis* (Drury, 1773)

462. *maesites* (Herrich-Schäffer, 1865)

463. *telea* (Hewitson, 1868)

Allosmaitia Clench, [1964]

464. *strophius* (Godart, [1824])

[*Allosmaitia pion* (Godman & Salvin 1887) is a synonym, *fide* Robbins (2004); also see Ferris (1989c: p. 28).]

Electrostrymon Clench, 1961*

465. *hugon* (Godart, [1824])

[Previously known as *E. endymion cyphara* (Hewitson, 1874), a name placed in synonymy by Robbins (2004). Also in 2004, Faynel and Balint reported that *sangala* (Hewitson, 1868) is a junior synonym of *hugon* (Godart, [1824]).]

466. *joya* (Dognin, 1895)

[Previously referred to as *E. canus* (Druce, 1907), a name placed in synonymy by

Robbins (2004).]

467. *angelia* (Hewitson, 1874)

Calycopis Scudder, 1876

468. *cecropis* (Fabricius, 1793)

469. *isobeon* (Butler & H. Druce, 1872)

Strymon Hübner, 1818

470. *melinus* (Hübner, 1818)

471. *avalona* (W.G. Wright, 1905)

472. *rufofusca* (Hewitson, 1877)

473. *albata* (C. Felder & R. Felder, 1865)

474. *alea* (Godman & Salvin, 1887)

475. *bebrycia* (Hewitson, 1868)

476. *yojoa* (Reakirt, [1867])

477. *cestri* (Reakirt, [1867])

478. *martialis* (Herrich-Schäffer, 1865)

479. *istapa* (Reakirt, [1867])

[Formerly known as *columella* (Fabricius, 1793), which was determined to be a Caribbean species by Robbins & Nicolay (1999).]

480. *bazochii* (Godart, [1824])

481. *acis* (Drury, 1773)

482. *limenia* (Hewitson, 1868)

483. *serapio* (Godman & Salvin, 1887)

Tmolus Hübner, [1819]

484. *echion* (Linnaeus, 1767)

Ministrymon Clench, 1961*

485. *leda* (W.H. Edwards, 1882)

486. *clytie* (W.H. Edwards, 1877)

487. *azia* (Hewitson, 1873)

[Opinion 2144 of the ICZN (2006) conserves *Thecla azia* Hewitson over the senior synonym *Thecla guacamagari* Wallengren, 1860.]

Strephonota K. Johnson, Austin, Le Crom & Salazar, 1997*

488. *tephraeus* (Geyer, 1837)

Oenomaus Hübner, [1819]

489. *ortygynus* (Cramer, 1779)

Parrhasius Hübner, [1819]

490. *m-album* (Boisduval & Leconte, 1833)

Hypostrymon Clench, 1961*

491. *critola* (Hewitson, 1874)

Erora Scudder, 1872

492. *laeta* (W.H. Edwards, 1862)

493. *quaderna* (Hewitson, 1868)

Tribe Polyommatini (Blues)

[Our generic arrangement of this tribe follows Bálint & Johnson (1995).]

Leptotes Scudder, 1876

494. *cassius* (Cramer, 1775)

495. *marina* (Reakirt, 1868)

Zizula Chapman, 1910

496. *cyna* (W.H. Edwards, 1881)

Brephidium Scudder, 1876

497. *exilis* (Boisduval, 1852)

498. *pseudofea* (Morrison, 1873)

[Brown & Heineman (1972), Riley (1975) and Smith et al. (1994) treat *pseudofea* as a separate species from *exilis* (based on genitalic differences between them reported by Comstock & Huntington (1943)), and treat *isophthalma* (Herrich-Schäffer, 1862) as a subspecies of *exilis*. Note that Bordelon & Knudson (2000) mention possible intermediates between *exilis* and *pseudofea* from Texas, and Pavulaan & Gatrell (1999) suggest *exilis* and *pseudofea* may be conspecific (as treated by Scott (1986)), but provide little evidence to support their claim; further study is badly needed.]

Cupido Schrank, 1801

499. *comyntas* (Godart, [1824])

500. *amyntula* (Boisduval, 1852)

[We follow Karsholt & Razowski (1996), Hesselbarth et al. (1995: 547), and Gorbunov (2001) in treating *Everes* Hübner, [1819] as a subgenus of *Cupido*.]

Celastrina Tutt, 1906

501. *ladon* (Cramer, 1780)

[A neotype of *Papilio ladon* Cramer from Anne Arundel County, Maryland was designated by Miller and Brown (1981). This species is considered by Wright (in litt.) to be endemic to the Appalachian and Ozarkian regions. The neotype, which may require redesignation to comply with Article 75 of the International Code of Zoological Nomenclature, possesses unique male sex scales not possessed by male of other taxa that were previously grouped with this species.]

502. *lucia* (W. Kirby, 1837)

[Evidence of this taxon being a full species is provided by Pavulaan (1995), Kondla (1999), Nielsen (1999), and Oehlenschlager & Huber (2002).]

503. *neglecta* (W.H. Edwards, 1862)

504. *echo* (W.H. Edwards, 1864)

[*Celastrina echo* was treated as a full species by Guppy & Shepard (2001).]

*[Further elaboration on the relationships between *echo*, *nigrescens* (Fletcher, 1903), *sidara* (Clench, 1944), *cinerea* (W.H. Edwards, 1883), *gozora* (Boisduval, 1870), and Baja California populations (see Brown et al. 1992) is needed to determine their status; some or all of which may represent species-level taxa.]

505. *nigra* (Forbes, 1960)*

[*Celastrina ebenina* Clench, 1972* is a synonym; see Scott & Wright (1993).]

506. *neglectamajor* Opler & Krizek, 1984*

507. *humulus* Scott & D. Wright, 1998*

508. *idella* Wright & Pavulaan, 1999*

508.1 *serotina* Pavulaan and Wright, 2005*

*[Additional species-level entities have been recognized by some (see Pratt et al. (1994), Wright (1995) and Layberry et al. (1998)), including *violacea* (W.H. Edwards, 1866).]

*[It is uncertain which, if any, of our North American *Celastrina* taxa are conspecific with the Palearctic *Celastrina argiolus* (Linnaeus, 1758), where they have traditionally been placed (see Eliot & Kawazoé 1983).]

Glauopsyche Scudder, 1872

509. *piasus* (Boisduval, 1852)

510. *lygdamus* (Doubleday, 1841)

[The extinct *G. xerxes* (Boisduval, 1852) was suggested to be a subspecies of *lygdamus* by Scott (1986).]

Philotes Scudder, 1876

511. *sonorensis* (C. Felder & R. Felder, 1865)

Philotiella Mattoni, [1978]

512. *speciosa* (Hy. Edwards, 1877)

513. *leona* Hammond & McCorkle, 2000*

Euphilotes Mattoni, [1978]

[Most recent treatments of this genus are by Mattoni (1989), Pratt (1988, 1994, 1999), Pratt & Ballmer (1993), and Pratt & Emmel (1998). It is likely that additional species-level taxa will be delineated in the future. Warren 2005 cites two putative undescribed species from Oregon in the *Euphilotes battoides* complex.]

514. *battoides* (Behr, 1867)

[Tentatively includes, among others, *centralis* (Barnes & McDunnough, 1917).]

515. *bernardino* (Barnes & McDunnough, 1916)

516. *glaucon* (W.H. Edwards, 1871)

[Warren tentatively includes *intermedia* (Barnes & McDunnough, 1917) and *oregonensis* (Barnes & McDunnough, 1917) and discusses variation of the putative species group in Oregon.]

517. *ellisi* (Shields, 1975)*

518. *baueri* (Shields, 1975)*

519. *enoptes* (Boisduval, 1852)

520. *ancilla* (Barnes & McDunnough, 1918)

520.1 *columbiae* (Mattoni, 1954)

[Warren 2005 raises taxon to species status and presents evidence based on distribution, hosts, and phenotype.]

521. *mojave* (Watson & W.P. Comstock, 1920)

522. *rita* (Barnes & McDunnough, 1916)

523. *pallescens* (Tilden & Downey, 1955)

524. *spaldingi* (Barnes & McDunnough, 1917)

Hemiargus Hübner, 1818

525. *ceraunus* (Fabricius, 1793)

[Note that Brown & Heineman (1972), Riley (1975), Smith et al. (1994) and Lamas (2004) treat *ceraunus* as a subspecies of *hanno* (Stoll, 1790); however, Clench (1977) and Johnson & Bálint (1995) consider the two as separate species, and Schwartz (1989) reported *hanno* and *ceraunus* as widely sympatric on Hispaniola. The situation needs further study.]

Echinargus Nabokov, 1945*

526. *isola* (Reakirt, [1867])

Cyclargus Nabokov, 1945*

527. *thomasi* (Clench, 1941)
528. *ammon* (Lucas, 1857)

[This species was first reported in our region by Beck (1985), from Big Pine Key, Monroe Co., Florida; it was subsequently photographed there by Krizek (1998).]

*[Generic treatment of *Hemiargus*, *Echinargus*, and *Cyclargus* follows Nabokov (1945), Smith et al. (1994), Bálint & Johnson (1995), Johnson & Bálint (1995), and Lamas (2004); also see Brown & Heineman (1972).]

Plebejus Kluk, 1780

529. *idas* (Linnaeus, 1761)
530. *anna* (W.H. Edwards, 1861)

[This taxon was treated as a full species by Bálint & Johnson (1997), Guppy & Shepard (2001), Pyle (2002), and Warren (2005); also see Nice & Shapiro (1999). More fieldwork and laboratory investigations are needed to clarify the situation in North America and Eurasia (see Tuzov et al. 2000).]

531. *mellissa* (W.H. Edwards, 1873)

[See Lane & Weller (1994) for details on the taxonomic status of *P. mellissa* populations, including *samuelis* (Nabokov, 1944).]

532. *saepiolus* (Boisduval, 1852)
533. *emigdionis* (F. Grinnell, 1905)
534. *icariooides* (Boisduval, 1852)
535. *shasta* (W.H. Edwards, 1862)
536. *acmon* (Westwood, [1851])
537. *lupini* (Boisduval, 1869)

[The previous two species appear to represent a species complex containing, at present, an unknown number of species. The group appears to be much more diverse than was suggested by Goodpasture (1973, 1974). Recently, Scott (1998) transferred several subspecies from *acmon* to *lupini*, but presented almost no data to support the changes. Warren (2005) discusses an undescribed species-level taxon from the Oregon Cascades. The complex is currently under study by PAO (Opler, 2003).]

538. *neurona* (Skinner, 1902)
539. *optilete* (Knoch, 1781)
540. *glandon* (de Prunner, 1798)
541. *podarce* (C. Felder & R. Felder, 1865)
542. *cassiope* (J. Emmel & T. Emmel, 1998)*

[Relationships of the above three taxa were discussed by Emmel & Emmel (1998), but are still under debate. Some authors recognize additional species, including *rusticus* (W.H. Edwards, 1865) and *franklinii* (Curtis, 1835); see Ferris (1989c).]

*[Our treatment of *Plebejus* follows that presented by Gorbunov (2001). He refers to it as a “super genus” with many subgenera, which in our area include *Lycaeides* Hübner, [1819], *Plebulina* Nabokov, [1945], *Icaricia* Nabokov, [1945], *Vacciniina* Tutt, 1909, and *Agriades* Hübner, [1819]. Bálint & Johnson (1997) present a considerably different arrangement of the group, placing our members in *Aricia* [Reichenbach], 1817, *Albulina* Tutt, 1909 and *Plebejus*. None of these authors retain

the use of *Lycaeides*, *Icaricia*, *Vacciniina* or *Plebulina* as valid genera.]

FAMILY RIODINIDAE

[Our treatment of the Riodinidae as a family follows Robbins (1988a,b) and Callaghan and Lamas (2004), but should be considered a tentative arrangement only. The relationship of this group to the Lycaenidae and Danainae requires further study, as noted by Campbell et al. (2000).]

Subfamily Riodininae

Calephelis Grote & Robinson, 1869

- 543. *virginiensis* (Guérin-Méneville, [1832])
- 544. *borealis* (Grote & Robinson, 1866)
- 545. *muticum* McAlpine, 1937
- 546. *nemesis* (W.H. Edwards, 1871)
- 547. *perditalis* Barnes & McDunnough, 1918
- 548. *wrighti* Holland, 1930
- 549. *rawsoni* McAlpine, 1939

[Scott (1986) lumped *freemani* McAlpine, 1971* and *arizonensis* McAlpine, 1971* as subspecies of *rawsoni*, yet presented no data in support of these changes (but did describe genitalic differences between these taxa). We do not accept his combinations, and stress that further study of this genus is badly needed; Neck (1996) treats *freemani* as a full species. The status of *C. dreisbachi* McAlpine, 1971* (one specimen from Arizona in the type series) needs clarification, and is tentatively not included in our list of species.]

- 550. *freemani* McAlpine, 1971*
- 551. *arizonensis* McAlpine, 1971*

Caria Hübner, 1823

- 552. *ino* Godman & Salvin, 1886

Lasaia H.W. Bates, 1868

- 553. *sula* Staudinger, 1888
- 554. *maria* Clench, 1972*

[Specimen collected in Guadalupe Canyon, Cochise Co., Arizona, by Kilian Roever; determination confirmed by R. Bailowitz.]

Melanis Hübner, [1819]

- 555. *pixe* (Boisduval, 1836)

Emesis Fabricius, 1807

- 556. *zela* Butler, 1870
- 557. *ares* (W.H. Edwards, 1882)

[Callaghan and Lamas (2004) treat *ares* as a synonym of *zela cleis* (W.H. Edwards, 1882), but we feel that further study of the types is needed, since two species indeed appear to be involved in our region (e.g. Bailowitz & Brock 1991).]

- 558. *emesia* (Hewitson, 1867)
- 559. *tenedia* C. Felder & R. Felder, 1861

Apodemia C. Felder & R. Felder, 1865

- 560. *mormo* (C. Felder & R. Felder, 1859)

561. *duryi* (W.H. Edwards, 1882)

[See Toliver et al. (1994) for notes on the specific status of *duryi*.]

562. *virgulti* (Behr, 1865)

563. *mejicanus* (Behr, 1865)

[Species-level relationships of the *mormo* complex (above four taxa) remain poorly understood, and our arrangement is tentative. Our treatment is based, in part, on information in Pratt & Ballmer (1991), Cary & Holland (1992) and Toliver et al.

(1994). The group has not been considered as a whole since the analysis of Opler & Powell (1962).]

564. *palmerii* (W.H. Edwards, 1870)

565. *hepburni* Godman & Salvin, 1886

566. *walkeri* Godman & Salvin, 1886

567. *multiplaga* Schaus, 1902

568. *phyciodoides* Barnes & Benjamin, 1924

[Possibly extirpated in U.S. portion of range (Bailowitz & Brock 1991); also see Holland & Forbes (1981).]

569. *nais* (W.H. Edwards, 1877)

570. *chisosensis* H.A. Freeman, 1964*

FAMILY NYMPHALIDAE

[Our subfamily and generic arrangement generally follows Harvey (1991), as modified by Ackery et al. (1999), Freitas (1999), Brower (2000), Wahlberg (2002), Freitas & Brown (in preparation) and Lamas (2004), although studies are under way to further resolve relationships. It is possible that the family, as presented here, is not a monophyletic group, since the exact placement of the Libytheinae and Danainae remains uncertain (see Campbell et al. 2000).]

Subfamily Libytheinae

Libytheana Michener, 1943

571. *carinenta* (Cramer, 1777)

[Many authors (e.g. Field 1940, Friedlander 1984, Shields 1984, 1985, Okano 1987, Ferris 1989c, Motono 1993, and Austin & Emmel 1998) treat *bachmani* (Kirtland, 1851) and *carinenta* as separate species. However, little information on populations across North and Central America is available, and reported genitalic differences between the two do not hold up in series (Kawahara 2001, pers. comm. 2002).]

572. *motya* (Hübner, [1823])

[Heitzman & Heitzman (1973) reported this species from Texas as rare strays or temporary colonists from Cuba; see Alayo and Hernández (1981).]

Subfamily Danainae

[In our area this subfamily includes the tribes Danaini and Ithomiini; see Ackery & Vane-Wright (1984), Ackery (1987), and Ackery et al. (1999).]

Danaus Kluk, 1780

573. *plexippus* (Linnaeus, 1758)

574. *gilippus* (Cramer, 1776)

575. *eresimus* (Cramer, 1777)

Lycorea Doubleday, [1847]

576. *halia* (Hübner, 1816)

[Central American populations are sometimes treated as species *cleobaea* (Godart, 1819), e.g. Miller & Brown (1981), Schwartz (1989), Brown (1992), and Smith et al. (1994); but see Lamas (2004).]

Dircenna Doubleday, 1847

577. *klugii* (Geyer, 1837)

Greta Heming, 1934

577.1 *morgane* (Geyer, 1837)

[First documented for the United States by Dauphin et al. 2005].]

Pteronymia Butler & Druce, 1872

577.2 *cotytto* (Guérin-Méneville, [1844])

[Bordelon and Knudson (2006) reported a female that was found and photographed at the NABA International Butterfly Park in Hidalgo County, Texas by Dauphin on December 7, 2005].

Subfamily Heliconiinae

Dione Hübner, [1819]

578. *moneta* Hübner, [1825]

Agraulis Boisduval & Leconte, [1835]

579. *vanillae* (Linnaeus, 1758)

Dryas Hübner, [1807]

580. *iulia* (Fabricius, 1775)

Dryadula Michener, 1942

581. *phaetusa* (Linnaeus, 1758)

Eueides Hübner, 1816

582. *isabella* (Stoll, 1781)

Heliconius Kluk, 1780

583. *charithonia* (Linnaeus, 1767)

[Spelling corrected to original orthography (Brower 1994).]

584. *erato* (Linnaeus, 1758)

Euptoieta Doubleday, 1848

585. *claudia* (Cramer, 1776)

586. *hegesia* (Cramer, 1779)

Speyeria Scudder, 1872

587. *diana* (Cramer, 1777)

588. *cybele* (Fabricius, 1775)

[Includes *leto* (Behr, 1862), although this is treated as a full species by some (e.g. Kondla 1999).]

589. *aphrodite* (Fabricius, 1787)

590. *idalia* (Drury, 1773)

591. *nokomis* (W.H. Edwards, 1862)

- 592. *edwardsii* (Reakirt, 1866)
- 593. *coronis* (Behr, 1864)
- 594. *zerene* (Boisduval, 1852)
- 595. *carolae* (dos Passos & Grey, 1942)

[Although somewhat intermediate between *coronis* and *zerene*, this isolated, disjunct taxon was considered a full species by Emmel & Austin (1998).]

- 596. *callippe* (Boisduval, 1852)
- 597. *egleis* (Behr, 1862)
- 598. *adiaste* (W.H. Edwards, 1864)
- 599. *atlantis* (W.H. Edwards, 1862)
- 600. *hesperis* (W.H. Edwards, 1864)

[Pyle (2002) treats the above two taxa as a single species based on unpublished research by Paul Hammond. Until this evidence is presented, we follow the arrangement proposed by Scott et al. (1998); also see Ferris (1983).]

- 601. *hydaspe* (Boisduval, 1869)
- 602. *mormonia* (Boisduval, 1869)

Boloria Moore, [1900]

[We follow a broad usage of *Boloria*; see dos Passos & Grey (1945), dos Passos (1964), Grey (1957, 1989) and Gorbunov (2001). *Clossiana* Reuss, 1920 and *Proclossiana* Reuss, 1926 are considered to be subgenera or synonyms (but see Warren (1944), Miller & Brown (1981) and Aubert et al. (1996).]

- 603. *alaskensis* (Holland, 1900)

[*Boloria napaea* (Hoffmansegg, 1804) is a separate Eurasian species with differing genitalia that reportedly occurs sympatrically with *alaskensis* at some locations. See discussions by Crosson du Cormier (1977), Dubatolov (1992), Kosterin (2000), Tuzov et al. (2000), and Gorbunov (2001).]

- 604. *eunomia* (Esper, 1800)
- 605. *selene* ([Denis & Schiffermüller], 1775)
- 606. *bellona* (Fabricius, 1775)
- 607. *frigga* (Thunberg, 1791)
- 608. *improba* (Butler, 1877)

[Includes *acrocnema* Gall & Sperling, 1980*.]

- 609. *kriemhild* (Strecker, 1879)
- 610. *epithore* (W.H. Edwards, 1864)
- 611. *polaris* (Boisduval, [1828])
- 612. *freija* (Thunberg, 1791)
- 613. *natazhati* (Gibson, 1920)

[Taxonomic status as a full species detailed by Troubridge & Wood (1995), Shepard et al. (1998), and Guppy & Shepard (2001).]

- 614. *alberta* (W.H. Edwards, 1890)
- 615. *astarte* (Doubleday, [1847])

[Tentatively includes *distincta* (Gibson, 1920), following dos Passos (1961); but see Wyatt (1958). The relationship of *astarte* to *distincta* and *tschukotkensis* (Wyatt, 1961) needs further elaboration. Guppy & Shepard (2001) very briefly summarize reasons for using *tritonia* (Böber, 1812) for our North American populations, a

combination also implied by Gorbunov (2000). However, no comprehensive review of the complex has been presented, and Tuzov et al. (2000) maintain *tritonia* and *distincta* as separate species.]

616. *chariclea* (Schneider, 1794)

[Includes our former concept of *titania* (Esper, [1793]), a wholly Palaearctic taxon (see Shepard 1998); also includes *montinus* (Scudder, 1863). However, the possibility exists that more than one species is being represented in our area by this name (see Layberry et al. 1998).]

Subfamily Nymphalinae

Poladryas Bauer, 1961

617. *minuta* (W.H. Edwards, 1861)

618. *arachne* (W.H. Edwards, 1869)

[We do not feel that Scott (1974, 1986) presents sufficient data to lump the above two taxa. Several authors also maintain *arachne* and *minuta* as full species, e.g. Ferris (1989c) and Austin (1998a); however, further research in the U.S. and Mexico is needed to resolve the situation.]

Chlosyne Butler, 1870

619. *theona* (Ménétriés, 1855)

[Includes *chinatiensis* (Tinkham, 1944), according to Austin & Smith (1998).]

620. *cyneas* (Godman & Salvin, 1878)

621. *fulvia* (W.H. Edwards, 1879)

622. *leanira* (C. Felder & R. Felder, 1860)

623. *californica* (W.G. Wright, 1905)

624. *lacinia* (Geyer, 1837)

625. *definita* (E.M. Aaron, [1885])

626. *endeis* (Godman & Salvin, 1894)

627. *janais* (Drury, 1782)

628. *rosita* A. Hall, 1924

629. *melitaeoides* (C. Felder & R. Felder, 1867)

[See Kendall & McGuire (1984) for reports from Texas.]

630. *eumeda* (Godman & Salvin, 1894)

[Kons (2000) and Luis et al. (2003) consider *eumeda*, *melitaeoides* and *marina* (Geyer, 1837) to be separate species (these were lumped by Scott (1986) without justification). The report from Pima Co., Arizona, appears to be of the semi-desert species, *eumeda*; however, *C. marina* also occurs in Sonora, Mexico, in montane habitats.]

631. *gorgone* (Hübner, [1810])

632. *nycteis* (Doubleday, [1847])

[A petition to the International Commission on Zoological Nomenclature to suppress *ismeria* (Boisduval & Leconte, [1835]), a possible senior synonym of *nycteis* (see Gatrell 1998, 2000), is in preparation; see ICZN (1999).]

633. *harrisii* (Scudder, 1864)

634. *palla* (Boisduval, 1852)

635. *gabbi* (Behr, 1863)

636. *acastus* (W.H. Edwards, 1874)

[Includes *neumoegeni* (Skinner, 1895), among others.]

637. *whitneyi* (Behr, 1863)

[Tentatively includes *damoetas* (Skinner, 1902), following Scott (1998) and Kons (2000), but further elaboration on the relationships of taxa in this group is needed.]

638. *hoffmanni* (Behr, 1863)

*[*Charidryas* Scudder, 1872 (see Higgins 1960) and *Thessalia* Scudder, 1875 (see Wahlberg & Zimmerman 2000) are considered synonyms of *Chlosyne*.]

Microtia H.W Bates, 1864

639. *elva* H.W. Bates, 1864

Dymasia Higgins, 1960*

640. *dymas* (W.H. Edwards, 1877)

Texola Higgins, 1959

641. *elada* (Hewitson, 1868)

Phyciodes Hübner, [1819]

642. *graphica* (R. Felder, 1869)

[*Phyciodes graphica*, described in April, 1869, supercedes *P. vesta* (W.H. Edwards, 1869), which was described in September-October, 1869; see Lamas (2004).]

643. *picta* (W.H. Edwards, 1865)

644. *orseis* W.H. Edwards, 1871

645. *pallida* (W.H. Edwards, 1864)

646. *mylitta* (W.H. Edwards, 1861)

647. *phaon* (W.H. Edwards, 1864)

648. *tharos* (Drury, 1773)

649. *cocyta* (Cramer, 1777)

[Supersedes *selenis* (W. Kirby, 1837), an apparent junior synonym; see Scott (1994).]

[Whether *Phyciodes incognitus* Gatrell, 2004 should be recognized as a separate species or referred to as a subspecies of *Phyciodes cocyta* remains to be ascertained (Gatrell 2004).]

650. *batesii* (Reakirt, [1866])

651. *pulchella* (Boisduval, 1852)

[Lectotype designation of *pulchella* by Emmel et al. (1998a) displaces *pratensis* (Behr, 1863). See Miller & Brown (1981) for synonymy of *campestris* (Behr, 1863).]

*[Our treatment of *Phyciodes* follows Wahlberg et al. (2003).]

Anthanassa Scudder, 1875

[Recognition of this genus follows Wahlberg & Zimmerman (2000).]

652. *frisia* (Poey, 1832)

653. *tulcis* (H.W. Bates, 1864)

654. *texana* (W.H. Edwards, 1863)

[*Anthanassa seminole* (Skinner, 1911) was suggested to be a species distinct from *texana* by Watts & Habeck (1991), and treated as such by Neck (1996); further study is needed to determine its status.]

655. *ptolyca* (H.W. Bates, 1864)

656. *argentea* (Godman & Salvin, 1882)

[First U.S. record reported by Chuah & Cushing (1995); also see Bordelon & Knudson (2000). *Anthanassa ardys* (Hewitson, 1864) may have also strayed into Texas (see Excluded Species).]

Tegosa Higgins, 1981*

657. *anieta* (Hewitson, 1864)

Euphydryas Scudder, 1872

658. *gillettii* (Barnes, 1897)

659. *phaeton* (Drury, 1773)

660. *editha* (Boisduval, 1852)

661. *chalcedona* (Doubleday, [1847])

662. *anicia* (Doubleday, [1847])

[We tentatively consider *anicia* to be a separate species from *chalcedona* following Ferris (1989b), Guppy & Shepard (2001) and Pyle (2002), however further elaboration on the relationship between the two taxa is needed. While the taxa clearly behave as two species in the Pacific Northwest, their relationship appears to be less well defined in other areas, e.g. Brussard et al. (1989), Austin & Murphy (1998). We tentatively consider *colon* (W.H. Edwards, 1881) to be conspecific with *chalcedona*, but again, much additional research is needed. A recent study by Zimmerman et al. (2000) did not resolve relationships among these taxa.]

*[Following Wahlberg & Zimmerman (2000) and Zimmerman et al. (2000), *Occidryas* Higgins, 1978* and *Hypodryas* Higgins, 1978* are considered synonyms of *Euphydryas*.]

Hypolimnas Hübner, [1819]

663. *misippus* (Linnaeus, 1764)

Junonia Hübner, [1819]

[Our use of *Junonia* instead of *Precis* Hübner, [1819] follows de Lesse (1952), Turner & Parnell (1985), and Lamas (2004); but see Hemming (1934) and Comstock (1944).]

664. *coenia* Hübner, [1822]

665. *evarete* (Cramer, 1779)

[Includes *nigrosuffusa* Barnes & McDunnough, 1916 (*contra* Turner & Parnell 1985), following Lamas (2004).]

666. *genoveva* (Stoll, 1780)

*[The relationships between these *Junonia* taxa need further elaboration; see Hafernik (1982), Turner & Parnell (1985), and Lamas (2004).]

Anartia Hübner, [1819]

667. *jatrophae* (Linnaeus, 1763)

668. *fatima* (Fabricius, 1793)

[Sometimes treated as a subspecies of *amathea* (Linnaeus, 1758), e.g. Lamas (2004); but see Silberglied et al. (1979), Dasmahapatra et al. (2002), and Blum et al. (2003).]

669. *chrysopelea* (Hübner, [1831])

[Reported as a subspecies of *lytrea* (Godart, 1819) by Anderson (1974), Bennett & Knudson (1976), Scott (1986), and Lamas (2004); but see Silberglied et al. (1979), Alayo & Hernández (1981), and Blum et al. (2003).]

Siproeta Hübner, [1823]

670. *stelenes* (Linnaeus, 1758)

671. *epaphus* (Latreille, [1813])

Polygonia Hübner, [1819]

672. *interrogationis* (Fabricius, 1798)

673. *comma* (Harris, 1842)

674. *satyrus* (W.H. Edwards, 1869)

675. *faunus* (W.H. Edwards, 1862)

676. *gracilis* (Grote & Robinson, 1867)

[Includes *zephyrus* (W.H. Edwards, 1870); see Scott (1984) and Layberry et al. (1998).]

677. *progne* (Cramer, 1775)

678. *oreas* (W.H. Edwards, 1869)

[This species was treated as distinct from *progne* (*contra* Scott 1984) by Bird et al. (1995), Layberry et al. (1998), and Guppy & Shepard (2001).]

Aglais Dalman, 1816

679. *milberti* (Godart, 1819)

680. *urticae* (Linnaeus, 1758)

[Several individuals of this common Palaearctic butterfly have been captured or photographed (e.g. Glassberg 1992, Zirlin & Ingraham 1997, Zirlin 2002), and this species is likely either an occasional accidental introduction or, less likely, a periodic vagrant.]

Nymphalis Kluk, 1780

681. *vaualbum* (Denis and Schiffermüller)

[Guppy & Shepard (2001: pp. 256-257) indicate that *vaualbum* may be a *nomen nudum* (see ICZN 1999), and use *l-album* (Esper, 1781) in its place. This treatment is repeated by Wahlberg & Nylin (2003). A petition to the ICZN is being prepared to resolve the issue, and until the commission rules, we maintain the name in widest use, *vaualbum*, as recommended by the ICZN (1999). See Sattler & Tremewan (1984) for further discussion.]

[The inclusion of *l-album* (formerly as *vau-album*) in *Roddia* Korshunov, 1995 by Guppy and Shepard (2001) and others is reversed by Wahlberg & Nylin (2003).]

682. *antiopa* (Linnaeus, 1758)

683. *californica* (Boisduval, 1852)

[Miller & Miller (1990) treat *californica* as a subspecies of the Eurasian *xanthomelas* Esper [1781], but see Nylin et al. (2001).]

*[Our arrangement of *Nymphalis* and *Aglais* follows Nylin et al. (2001) as modified by Wahlberg and Nylin (2003).]

Vanessa Fabricius, 1807

684. *atalanta* (Linnaeus, 1758)

685. *cardui* (Linnaeus, 1758)

686. *annabella* (Field, 1971)*

687. *virginiensis* (Drury, 1773)

[Some authors place the above three species in *Cynthia* Fabricius, 1807, following Field (1971), but several natural hybrids between *atalanta* and *annabella* are known (Comstock 1927, Dimock 1973, Tilden & Smith 1986).]

Hypenanartia Hübner, [1821]

688. *lethe* (Fabricius, 1793)

Historis Hübner, [1819]

689. *odius* (Fabricius, 1775)

690. *acheronta* (Fabricius, 1775)

Smyrna Hübner, [1823]

691. *blomfildia* (Fabricius, 1781)

Subfamily Limenitidinae

[Our treatment of the Limenitidinae and Biblidinae follows Freitas (1999), Wahlberg (2002) and Freitas & Brown (in preparation), although Lamas (2004) places this subfamily within the Biblidinae, and *Marpesia* may belong elsewhere; further study is needed.]

Limenitis Fabricius, 1807

[*Basilarchia* Scudder, 1872 is considered a synonym; see Chermock (1950), Niculescu (1986), Tuzov et al. (2000), and Gorbunov (2001).]

692. *arthemis* (Drury, 1773)

693. *archippus* (Cramer, 1776)

694. *lorquini* (Boisduval, 1852)

695. *weidemeyerii* W.H. Edwards, 1861

[Suggested to be conspecific with *lorquini* by Porter (1990), but see Boyd et al. (1999).]

Adelpha Hübner, [1819]

696. *bredowii* Geyer, 1837

697. *fessonia* (Hewitson, 1847)

698. *basiloides* (H.W. Bates, 1865)

[Opler (1992) and Neck (1996) reported this species from Texas.]

Marpesia Hübner, 1818

699. *chiron* (Fabricius, 1775)

700. *petreus* (Cramer, 1776)

701. *eleuchea* Hübner, 1818

702. *zerynthia* Hübner, [1823]

[This species is occasionally listed as *M. coresia* (Godart, [1824]), a junior synonym of *zerynthia* (see Neild 1996).]

Subfamily Biblidinae

Biblis Fabricius, 1807

703. *hyperia* (Cramer, 1779)

Mestra Hübner, [1825]

704. *amymone* (Ménétriés, 1857)

[This is sometimes treated as a subspecies of the Jamaican *M. dorcus* (Fabricius, 1775), e.g. Lamas (2004).]

Eunica Hübner, [1819]

705. *monima* (Stoll, 1782)

706. *tatila* (Herrich-Schäffer, [1855])

Myscelia Doubleday, [1845]

707. *ethusa* (Doyère, [1840])

708. *cyananthe* C. Felder & R. Felder, 1867

Dynamine Hübner, [1819]

709. *dyonis* Geyer, 1837

[Sometimes considered to be a subspecies of the Caribbean *D. egaea* (Fabricius, 1775), e.g. de la Maza & Turrent (1985) and Smith et al. (1994), but *serina* (Fabricius, 1775) has precedence over *egaea* (see Lamas 2004).]

709.1. *postverta* (Cramer, 1779)

[Bordelon & Knudson (2006) reported that three individuals of *D. postverta mexicana* d'Almeida were found in Hidalgo County, Texas during December, 2005. These butterflies were photographed by a number of observers and images of two of the individuals were illustrated by Bordelon and Knudson. Previously, the species was reported from Texas by Pyle (1986) as *D. mylitta* (Cramer, 1779) and by Stanford (2002), but there was no satisfactory documentation.]

Diaethria Billberg, 1820

710. species

[The west Texas *Diaethria* record (see Kendall & McGuire 1984, Neck 1996) is apparently of *D. anna* (Guérin-Méneville, [1844]) or *D. astala* (Guérin-Méneville, [1844]), since these are the only *Diaethria* species resident in northeastern Mexico (de la Maza & Turrent 1985); however, the west Texas specimen has not been positively determined and is reportedly no longer extant. There are also unverified records for both *anna* and *astala* from south Texas (Stanford 2002). See Excluded Species.]

Epiphile Doubleday, [1845]

711. *adrasta* Hewitson, 1861

Temenis Hübner, [1819]

711.1 *laothoe* (Cramer, [1777])

[First record from United States reported by Grishin (2005).]

Hamadryas Hübner, [1806]

712. *februa* (Hübner, [1823])

713. *amphichloe* (Boisduval, 1870)

714. *glaucome* (H.W. Bates, 1864)

715. *atlantis* (H.W. Bates, 1864)

716. *feronia* (Linnaeus, 1758)

717. *guatemalena* (H.W. Bates, 1864)

718. *iphthime* (H.W. Bates, 1864)

719. *amphinome* (Linnaeus, 1767)

Subfamily Charaxinae

Anaea Hübner, [1819]

720. *troglodyta* (Fabricius, 1775)

[Includes *floridalis* F. Johnson & W.P. Comstock, 1941, in accordance with Lamas (2004), but see discussion by Smith et al. (1994: p. 65).]

721. *aidea* (Guérin-Méneville, [1844])

[Often considered a species-level taxon (e.g. DeVries 1987), but sometimes treated as a subspecies of *troglodyta* (e.g. Lamas 2004).]

722. *andria* Scudder, 1875

Memphis Hübner, [1819]

723. *glycerium* (Doubleday, [1849])

[Some authors (e.g. Lamas 2004) place *glycerium* in the genus *Fountainea* Rydon, 1971*, but the monophyly of *Fountainea* has not been demonstrated and is questionable.]

724. *pithyusa* (R. Felder, 1869)

725. *echemus* (Doubleday, [1849])

Subfamily Apaturinae

Asterocampa Röber, 1915

726. *celtis* (Boisduval & Leconte, [1835])

727. *leilia* (W.H. Edwards, 1874)

728. *clyton* (Boisduval & Leconte, [1835])

729. *idyja* (Geyer, [1828])

[It is possible that *Asterocampa argus* (H.W. Bates, 1864), currently treated as the continental subspecies of *idyja*, may comprise a separate species (see Smith et al. 1994).]

*[Our treatment of *Asterocampa* species follows Friedlander (1987), although further study of the genus is needed.]

Doxocopa Hübner, [1819]

730. *laure* (Drury, 1773)

731. *pavon* (Latreille, [1809])

Subfamily Morphinae

Morpho Fabricius, 1807

732. *polyphemus* Westwood, [1850]

Subfamily Satyrinae

[Our arrangement of this subfamily generally follows the classification presented by Miller (1968), as modified by Harvey (1991).]

Enodia Hübner, [1819]

733. *portlandia* (Fabricius, 1781)

734. *anthedon* A.H. Clark, 1936

735. *creola* (Skinner, 1897)

Satyrodes Scudder, 1875

736. *eurydice* (Linnaeus, 1763)

737. *appalachia* (R.L. Chermock, 1947)

Cyllopsis R. Felder, 1869

738. *pyracmon* (Butler, 1867)

[*Cyllopsis henshawi* (W.H. Edwards, 1876) is treated as a seasonal form of *pyracmon*, as suggested by Bailowitz & Brock (1991) and Brock (1998); but see

Miller (1974).]

739. *pertepida* (Dyar, 1912)

740. *gemma* (Hübner, 1808)

Hermeuptychia Forster, 1964*

741. *sosybius* (Fabricius, 1793)

[The status of *sosybius* versus *hermes* (Fabricius, 1775) is uncertain and requires detailed study; see Forster (1964), Miller & Brown (1981: p. 241, note 624), and Smith et al. (1994). The use of *hermes* prevails in the Neotropical literature, e.g. Lamas (2004), for what may be (at least in part) the same species that occurs in our region.]

Neonympha Hübner, [1818]

742. *areolatus* (J.E. Smith, 1797)

743. *helicta* (Hübner, 1808)

[Species status recently proposed by Gatrell (1999); however, further research on this group is needed to clarify overall distributions and confirm the status of *helicta*.]

744. *mitchellii* French, 1889

Megisto Hübner, [1819]

745. *cymela* (Cramer, 1777)

[Includes *viola* (Maynard, 1891), following Catling & Calhoun (1997); however, it is still possible that more than one species is included under the name *cymela*, and further study is needed.]

746. *rubricata* (W.H. Edwards, 1871)

Paramacera Butler, 1868

747. *xicaque* (Reakirt, [1867])

[Tentatively includes *allyni* L. Miller, 1972*, following Scott (1986), but see Miller (1972). Further study is needed.]

Coenonympha Hübner, [1819]

748. *haydenii* (W.H. Edwards, 1872)

749. *tullia* (Müller, 1764)

[Treated as a species complex by Ferris (1989c); also see Davenport (1941) and Brown (1955, 1961). Included here are several taxa sometimes considered full species (e.g. dos Passos 1958); these include (among others) *ampelos* W.H. Edwards, 1871, *california* Westwood, 1851, *inornata* W.H. Edwards, 1861, *kodiak* W.H. Edwards, 1869, *ochracea* W.H. Edwards, 1861, and *nipisiquit* McDunnough, 1939. Layberry et al. (1998) treated *nipisiquit* as a full species based on (then) unpublished evidence by R. Webster; Webster (1999) subsequently treated *nipisiquit* as a subspecies of *tullia* but stressed that additional research was needed to determine its status. Handfield (1999) gave *nipisiquit* species-level status. Although gene flow between purported species was demonstrated by Porter & Geiger (1988) (also see Porter & Mattoon 1989), it is unclear which, if any, of our taxa are conspecific with the Eurasian *C. tullia*; see Kondla (1999) and Guppy & Shepard (2001). It is likely that more than one species-level taxon is included in this complex. Further research is badly needed since there is currently no satisfactory taxonomic arrangement for this group.]

Cercyonis Scudder, 1875

- 750. *pegala* (Fabricius, 1775)
- 751. *sthenele* (Boisduval, 1852)
- 752. *meadii* (W.H. Edwards, 1872)
- 753. *oetus* (Boisduval, 1869)

Erebia Dalman, 1816

- 754. *vidleri* Elwes, 1898
- 755. *rossii* (Curtis, 1835)
- 756. *disa* (Thunberg, 1791)
- 757. *mancinus* Doubleday, [1849]
[See Layberry et al. (1998) for separation of *mancinus* from *disa*.]
- 758. *magdalena* Strecker, 1880
- 759. *mackinleyensis* Gunder, 1932
[See Hilchie (1990) and Layberry et al. (1998) for separation of *mackinleyensis* from *magdalena*; however, Gorbunov (2001) treats the two as conspecific.]
- 760. *fasciata* Butler, 1868
- 761. *discoidalis* (W. Kirby, 1837)
- 762. *pawloskii* Ménétriés, 1859

[North American populations of this species have been referred to as *Erebia theano* (Tauscher, 1809), but Guppy & Shepard (2001) note that *Erebia theano* is restricted to eastern Siberia and Mongolia. See Tuzov et al. (1997). Gorbunov (2001, plate 41) illustrates the genitalia of both species.]

- 763. *youngi* Holland, 1900
- 764. *occulta* Roos & Kimmich, 1983*

[Our use of *occulta* over *anyuica* Kurentzov, 1966 follows Roos & Arnscheid (1984), Tuzov et al. (1997) and Belik & Zamolodchikov (2002); but see Dubatolov (1992) and Layberry et al. (1998).]

- 765. *lafontainei* Troubridge & Philip, 1983*
- 766. *callias* W.H. Edwards, 1871
- 767. *episodea* Butler, 1868

Gyrocheilus Butler, 1867

- 768. *patrobas* (Hewitson, 1862)

Neominois Scudder, 1875

- 769. *ridingsii* (W.H. Edwards, 1865)

[Scott (1998) described “*Hipparchia (Neominois) ridingsii wyomingo*” as a “new subspecies (or species?),” and suggested it is genetically isolated from *ridingsii*; Opler (1999) subsequently treated this as a full species. Additional field observations have shed some light on the situation but many questions remain and the status of *wyomingo* remains uncertain. The situation is currently under study by Matthew Garhart.]

Oeneis Hübner, [1819]

- 770. *philipi* Troubridge, 1988*

[The name *rosovi* Kurentzov, 1970 (a potential senior synonym of *philipi*) cannot be used in any meaningful way until a lectotype is designated, since two syntypes exist (but see Layberry et al. 1998). *Oeneis rosovi* was treated as a subspecies of *O. norna*]

(Thunberg, 1791) by Lukhtanov (1989), and as a synonym of that species by Gorbunov (2001). See also Lukhtanov & Eitschberger (2000).]

771. *polixenes* (Fabricius, 1775)

772. *jutta* (Hübner, [1806])

773. *mellissa* (Fabricius, 1775)

774. *alpina* Kurentzov, 1970

[Includes *excubitor* Troubridge, Philip, Scott & Shepard, 1982* as a synonym, following Layberry et al. (1998).]

775. *bore* (Esper, 1798)

[Tentatively includes *taygete* Geyer, [1830], following Layberry et al. (1998), but further study is needed to clarify the relationship of these taxa; Tuzov et al. (1997) consider the taxa to be separate species.]

776. *chryxus* (Doubleday, [1849])

[Includes *ivallda* (Mead, 1878), following Porter & Shapiro (1991).]

777. *alberta* Elwes, 1893

778. *nevadensis* (C. Felder & R. Felder, 1867)

779. *macounii* (W.H. Edwards, 1885)

780. *uhleri* (Reakirt, 1866)

[Hassler & Feil (2002) recently reported the occurrence of *Oeneis nanna* (Ménétriés, 1859) in North America, with the description of a new subspecies (*kluanensis*). The authors did not examine sufficient material of *uhleri*, and the proper taxonomic placement of this taxon is currently being studied.]

Hawaiian Butterflies

Although not part of the North American continent, Hawaii is one of the 50 United States. For sake of completeness we present a list of the butterflies recorded from the Hawaiian Islands, based on Riotte & Uchida (1979). Additional information is provided by Zimmerman (1958).

Family Hesperiidae

Erionota torus Evans, 1941

[Originally misdetermined as *E. thrax* (Linnaeus, 1767) by Riotte & Uchida (1979).]

Hylephila phyleus (Drury, 1773)

Family Papilionidae

Papilio xuthus Linnaeus, 1767

Family Pieridae

Pieris rapae (Linnaeus, 1758)

Phoebe agarithe (Boisduval, 1836)

Family Lycaenidae

Tmolus echion (Linnaeus, 1767)

Strymon bazochii (Godart, [1824])

Lampides boeticus (Linnaeus, 1767)

Brephidium exilis (Boisduval, 1852)

**Udara blackburni* (Tueyl, 1878)

[See Eliot & Kawazoé (1983) for generic combination.]

Family Nymphalidae

Agraulis vanillae (Linnaeus, 1758)

Vanessa virginiensis (Drury, 1773)

Vanessa cardui (Linnaeus, 1758)

Vanessa atalanta (Linnaeus, 1758)

**Vanessa tameamea* Eschscholtz, 1821

Danaus plexippus (Linnaeus, 1758)

*Endemic to Hawaiian Islands. All others are introduced exotics.

Excluded Species

The species listed below have been excluded from the North American fauna due to erroneous or inadequate documentation, or recent change in status.¹ This is not an exhaustive list, but treats most discrepancies in the post-1980 literature that are not discussed in the text. See Calhoun (1997) for a complete list of erroneous Florida records.

Hesperiidae

Phocides urania (Westwood, [1852])

[Dubious Texas records reported by Aaron (1890); Arizona records are unsubstantiated (see Bailowitz & Brock 1991).]

Polythrix asine (Hewitson, 1867)

[Erroneously reported from the U.S. based on an unjustifiable assumption of conspecificity with *mexicanus* by Scott (1986); see Freeman (1969) and Austin & Warren (2002).]

Polythrix procerus (Plötz, 1880)

[Misidentified but later corrected to *P. octomaculata* by Freeman (1967); *procerus* is now placed in *Cephise* Evans, 1952* (see Austin & Mielke 2000).]

Codatractus melon (Godman & Salvin, 1893) [Unsubstantiated records; see Bailowitz & Brock (1991: p. 25).]

Urbanus evona Evans, 1952

[Unsubstantiated sight record (Glassberg 2004). Genitalic examination is necessary for definite identification (Evans 1952, Warren, personal communication).]

Urbanus pronta Evans, 1952*

[Record refers to *U. pronus*; see Kendall & McGuire (1984).]

Urbanus albimargo (Mabille, 1876)

[Unsubstantiated records from Texas apparently refer to *doryssus*; see Ferris (1989c: p. 6).]

Astraptes galesus (Mabille, 1888)

[Unsubstantiated records; see Bailowitz & Brock (1991: pp. 30-31). The species is unknown from Mexico (Warren 2000, 2002).]

Thorybes valeriana (Plotz, 1881)

[Our records refer to *Codatractus mysie* (see Burns 1996); however the identity of *valeriana* remains a mystery.]

Dyscophellus euribates (Stoll, 1782)

[The report of this species from Texas by Aaron (1890), as *Eudamus hesus* (Westwood, [1852]), is unsubstantiated (see Lindsey et al. 1931); *euribates* remains unknown from Mexico (Warren 2000, 2002).]

Cogia cajeta (Herrich-Schäffer, 1869)

[One unsubstantiated record from Texas (Stanford 2002).]

Pellicia costimacula Herrich-Schäffer, 1870

[This South American species was shown to be separate from *arina* by Steinhauser (1989).]

Pellicia angra Evans, 1953*

[All reports from our area (e.g. Tilden 1974) apparently refer to *Pellicia arina*.]

Staphylus azteca (Scudder, 1872)

[Single individual reported from Texas by Freeman (1977) later identified as *Staphylus ceos* by ADW, not *mazans* as reported by Cassie et al. (2001).]

Pyrgus adepta Plötz, 1884

[Two highly unlikely and unverified records exist from Texas (Stanford 2002).]

Heliopyrgus sublinea (Schaus, 1902)

[Unsubstantiated record for this species from Texas reported by Stanford (2002); see Austin & Warren (2001) for generic combination.]

Piruna cyclosticta (Dyar, 1920)

[The recent report of *Piruna pirus* from Jeff Davis Co., Texas (Bordelon 2000), may represent this species, as listed (with a "?") by Stanford (2002). We have not yet examined these specimens.]

Callimormus saturnus (Herrich-Schäffer, 1869)

[A single male of this species labeled from Texas (Evans 1955) is most likely mislabeled, although the species is common in tropical northern Mexico.]

Repens florus (Godman, 1900)

[A single unverified record exists from south Texas (Stanford 2002), although this species is resident in Sonora and Tamaulipas, Mexico; see Vargas et al. (1996), Warren et al. (1998) and Warren (2000, 2002) for generic combination.]

Remella remus (Fabricius, 1798)

[A single unsubstantiated record exists from Texas (Stanford 2002).]

Decinea huasteca (H.A. Freeman, 1969)*

[Although cited by Ferris (1989c), no U.S. voucher is known.]

Polites subreticulata (Plötz, 1883)

[Reports by Freeman (1951) of this species apparently refer to *P. carus*; see Burns (1994a).]

Anatrytone potosiensis (H.A. Freeman, 1969)*

[Although cited by Ferris (1989c), no U.S. specimens are known; see Burns (1994b) for generic combination.]

Choranthus radians (Lucas, 1857)

[Unsubstantiated records from Florida; see Smith et al. (1994) and Calhoun (1997).]

Choranthus haitensis Skinner, 1920

[Unsubstantiated records from Florida; see Smith et al. (1994).]

Choranthus vitellius (Fabricius, 1793)

[Single female of this species reportedly from Florida (Evans 1955) is most likely mislabeled; see Smith et al. (1994).]

Quasimellana mexicana (Bell, 1942)

[Cited by Miller & Brown (1981) and Ferris (1989c), but apparently no known U.S. specimen; see Bailowitz & Brock (1991) and Burns (1994b).]

Atrytonopsis ovinia (Hewitson, 1866)

[Burns (1983) treats *A. ovinia* and *A. edwardsi* as separate species.]

Amblyscirtes fluonia Godman, 1900

[Reports of this species from Texas (Stanford 2002) are unsubstantiated.]

Panoquina fusina (Hewitson, 1868)

[Erroneously reported from North America based on an assumption of conspecificity with *evansi* by Evans (1955) and Scott (1986).]

Agathymus remingtoni (D. Stallings & Turner, 1958)

[Confusion with *estelleae* based on Scott (1986) has led to erroneous reports of this species in the U.S.]

Stallingsia smithi (H. Druce, 1896)

[This was erroneously reported from the U.S. based on an assumption of conspecificity with *maculosus* by Scott (1986).]

Papilionidae

Battus devilliers (Godart, [1824])

[Unsubstantiated records from Florida; see Smith et al. (1994).]

Eurytides celadon (Lucas, 1852)

[Considered hypothetical for Florida; see dos Passos (1961) and Smith et al. (1994).]

Papilio kahli F. Chermock & R. Chermock, 1937

[Excluded as hybrid on basis of Layberry et al. (1998); also see Klassen et al. (1989).]

Pieridae

Enantia mazai Llorente, 1984

[Report of this species by Stanford (2002) from Texas apparently refers to *E. albania*; see Kendall (1974a), Llorente (1984) and Llorente et al. (1997).]

Pontia callidice (Hübner, [1800])

[Reports of *callidice* from our area refer to *P. occidentalis*; the two were considered conspecific by Higgins & Riley (1970) and Scott (1986). Also see Shapiro (1976) and Chew & Watt (2006).]

Pontia chloridice (Hübner, [1813])

[Erroneously reported from our area by Scott (1986), based on an unsupported assumption of conspecificity with *beckerii*. Chew & Watt (2006) provide evidence of its distinctness.]

Pieris napi (Linnaeus, 1758)

[A Palaearctic species; see Chew & Watt (2006) and text.]

Pieris brassicae (Linnaeus, 1758)

[Accidental introductions, but no known persistent North American colonies (see Mello 1999 [these are sight records], Cassie et al. 2001, and Zirlin 2002).]

Anthocharis dammersi J.A. Comstock, 1929

[Considered to be a hybrid on basis of Emmel & Emmel (1973: p. 24) and Shields & Mori (1979).]

Phoebis intermedia (Butler, 1872)

[Reports of this taxon (e.g. Scott 1986) and of *P. rurina* (C. Felder & R. Felder, 1861) from our area refer to *P. neocypris*; *intermedia* is a synonym of *neocypris virgo* (Butler, 1870) (see Lamas 2004).]

Pyrisitia chamberlaini (Butler, 1898)

[Unsubstantiated report from Florida; see Scott (1986) and Smith et al. (1994).]

Lycaenidae

Eumaeus minyas (Hübner, [1809])

[Misidentification of *E. toxœus*; see Kendall & McGuire (1984).]

Pseudolycaena marsyas (Linnaeus, 1758)

[Dubiously reported from our region by Pyle (1981); *Pseudolycaena damo* (H. Druce, 1875) is the continental species which remains a hypothetical stray for Texas.]

Cyanophrys amyntor (Cramer, 1775)

[Specimen from Brewster County, Texas, reported as this species, was misidentified. Its true identity as *C. herodotus* confirmed by Johnson & Le Crom (1997: p. 26).]

Cyanophrys longula (Hewitson, 1868)

[Reported from southeastern Arizona by unsubstantiated sight records (Bailowitz & Brock 1991: p. 182), and one unverified record exists for Texas (Stanford 2002); no specimen or photographic voucher has been located.]

Rekoa zebina (Hewitson, 1869)

[Excluded from the North American fauna by Robbins (1991); records refer to misdetermined *R. marius*.]

Riodinidae

Euselasia abreas (W.H. Edwards, 1881)

[Described from Arizona, but likely based on mislabeled material (Powell 1975); it is unknown from Mexico (de la Maza et al. 1989) or Costa Rica (DeVries 1997), and its identity was unknown to D'Abrera (1994). However, this taxon may be a senior synonym of *E. sergia* (Godman & Salvin, 1885); see Lamas (2004).]

Calephelis nilus (C. Felder & R. Felder, 1861)

[Erroneously reported from our area by Scott (1986); records refer to *perditalis* (see McAlpine 1971).]

Calephelis sinaloensis McAlpine, 1971*

[Reported occurrence in Texas by Durden (1982) requires confirmation.]

Caria dominianus (Fabricius, 1793)

[Casually reported from Texas by DeVries (1997), but no details on this record have been presented.]

Nymphalidae

Greta polissena (Hewitson, 1863)

[Highly questionable records from Texas summarized by Kendall & McGuire (1984: pp. 41-41); the species is unknown from Mexico (de la Maza et al. 1989).]

Philaethria diatonica (Fruhstorfer, 1912)

[Sight record only, from Texas (as *dido*); see Bordelon (1991). *Philaethria diatonica* is considered to be a separate species from *dido* (Linnaeus, 1763) by Lamas (2004).]

Boloria titania (Esper, [1793])

[According to Shepard (1998), this is a strictly Palaearctic species.]

Chlosyne ehrenbergii (Geyer, [1833])

[Highly questionable record from Texas summarized by Kendall & McGuire (1984: p. 26).]

Chlosyne marina (Geyer, 1837)

[This was erroneously reported from the U.S. based on an assumption of conspecificity

with *melitaeoides* and *eumeda* by Scott (1986). See text.]

Chlosyne erodyle (H.W. Bates, 1864)

[Casually reported from Texas by Higgins (1960), but no authentic voucher specimen is known; see Neck (1996).]

Anthanassa ardys (Hewitson, 1864)

[While the single record of this species from Texas may be valid (see Stanford & Opler 1993), details have not been presented in the literature and we have not examined the specimen.]

Anthanassa drusilla (C. Felder & R. Felder, 1861)

[A single unsubstantiated record for this species exists from south Texas (Stanford 2002).]

Inachis io (Linnaeus, 1758)

[Three or four almost certainly human-related, accidental introductions are known; see Hinchliff (1994: p. 176), Anonymous (1995), and Zirlin (2002).]

Vanessa carye (Hübner, [1812])

[This South American species is occasionally reported from our area instead of *annabella* (e.g. Scott 1986); but see Field (1971) and Shapiro & Geiger (1989).]

Smyrna karwinskii Geyer, [1833]

[All records from North America apparently represent misidentifications of *S. blomfildia* (Kendall & McGuire 1984: p. 36-37).]

Mestra cana (Erichson, [1849])

[Reported as *cana floridana* Strecker, 1900, but it is likely that the types were mislabeled specimens; see Kimball (1965) and Masters (1970). Lamas (2004) treats *cana* and *floridana* as synonyms of *dorcas hersilia* (Fabricius, 1776).]

Dynamine tithia (Hübner, 1823)

[Texas specimen correctly identified and illustrated by Kendall & McGuire (1984: pp. 31-32); however it is a very unlikely long-distance stray and it may have been accidentally transported. The species is unknown from Mexico (de la Maza & Turrent 1985, de la Maza et al. 1989). The specimen is in Illinois Natural History Survey.]

Diaethria asteria (Godman & Salvin, 1894)

[Summary of history surrounding the single specimen reported from Texas provided by Kendall & McGuire (1984: p. 33). This record was seriously doubted by de la Maza & Turrent (1985: p. 32), and shown to be virtually impossible by Luis et al. (1996).]

Diaethria clymena (Cramer, 1775)

[Some records of *D. clymena* from our area, e.g. Klots (1951), apparently refer to this taxon; however, *clymena* is unknown from Mexico or the Caribbean (Kendall & McGuire 1984, de la Maza & Turrent 1985, Smith et al. 1994), and its occurrence in our area is assumed to be accidental.]

Hamadryas fornax (Hübner, [1823])

[Records of this species from the U.S. are based on misdetermined specimens (Jenkins 1983); but see Stanford (2002).]

Morpho peleides Kollar, 1850

[Texas sighting reported by Stallings & Turner (1946), Freeman (1960), and Neck (1996: p. 162.); see Penz & DeVries (2002) for systematics of the genus *Morpho*.]

Opsiphanes boisduvallii Doubleday, [1849]

[Circumstances surrounding a specimen of uncertain origin are discussed by Cassie et al. (2001).]

Erebia dabanensis Ershov, 1871

[Erroneously reported from North America based on an unsupported assumption of conspecificity with *youngi* by Scott (1986); see Troubridge & Philip (1983), Tuzov et al. (1997) and Belik & Zamolodchikov (2002).]

Erebia kozhantshikovi Sheljuzhko, 1925

[Erroneously reported from North America based on an unsupported assumption of conspecificity with *lafontainei* by Scott (1986); see Troubridge & Philip (1983) and Tuzov et al. (1997).]

Erebia inuitica Wyatt, 1966*[Known from only a single specimen whose authenticity is dubious; see Warren (1968) and Ferris (1989c: p. 59).]

Erebia theano (Tauscher, 1909)

[Although North American populations of this species have been referred to as *Erebia theano* (Tauscher, 1809) through several centuries, recently Guppy & Shepard (2001) demonstrated that *Erebia theano* is restricted to eastern Siberia and Mongolia. See Tuzov et al. (1997) for illustration of adult Eurasian *theano* and *pawloskii* Ménétriés, 1859, the latter which represents our North American butterfly. Gorbunov (2001, plate 41) illustrates the genitalia of both species. All North American infraspecific names now fall under *pawloskii*.]

\1 Adequate documentation should consist of a specimen or photograph (when genitalic dissection is not necessary), identified by a specialist in the appropriate taxonomic group. Details should be reported in the literature (website or e-mail postings are not adequate), along with complete data, when possible. Preferably, it should be deposited in a major entomological institution [National Museum of Natural History, Washington, D.C.; American Museum of Natural History, New York; Natural History Museum of Los Angeles County; California Academy of Sciences, San Francisco; Allyn Museum of Entomology, Sarasota; Carnegie Museum of Natural History, Pittsburgh; Canadian National Collection, Ottawa, Ontario] or university. To our knowledge no major collections have established photographic documentation files, but they should. Sight records do not constitute adequate documentation for national records. Moreover, photographs of living individuals of species that may be identified only by dissection do not constitute adequate documentation.

References

- Aaron, E.M. 1890. North American Hesperiidae. Entomological News 1(2):23-26.
- Ackery, P.R. 1987. The Danaid genus *Tellervo* (Lepidoptera, Nymphalidae)- a cladistic approach. Zoological Journal of the Linnean Society 89(3):203-274.
- Ackery, P.R., R. de Jong, & R.I. Vane-Wright. 1999. The butterflies: Hedyloidea, Hesperioidea, and Papilioidea. Pp. 263-300 in: Kristensen, N.P., editor. Lepidoptera, Moths and Butterflies. 1. Evolution, Systematics, and Biology. Handbuch der Zoologie. 4(35):263-300. Walter de Gruyter, Berlin. x + 491 pp.
- Ackery, P.R. & R.I. Vane-Wright. 1984. Milkweed Butterflies: Their Cladistics and Biology. Cornell University Press, Ithaca, New York. ix + 425 pp.
- Adams, J.K. 1984. An old first United States record finally published: *Papilio victorinus* (Papilionidae) in Laredo, Texas. Journal of the Lepidopterists' Society 37(4):3-18.
- Alayo, P.D. & L.R. Hernández. 1981. Atlas de las Mariposas Diurnas de Cuba (Lepidoptera: Rhopalocera). Editorial Científico-Técnica, Havana, Cuba. [ii] + 148pp., 49 pls.
- Anderson, R.A. 1974. Three new United States records (Lycaenidae and Nymphalidae) and other unusual captures from the lower Florida Keys. Journal of the Lepidopterists' Society 28(4):354-358.
- Anonymous. 1995. Reports of “wow!” status. American Butterflies 3(4):31.
- Anonymous. 2000. Run, don’t walk, for stray adventures. American Butterflies 8(1):43.
- Aubert, J., B. Barascud, H. Descimon, & F. Michel. 1996. Systématique moléculaire des argynnes (Lepidoptera, Nymphalidae). Comptes Rendus de l’Academie des Sciences Serie III, Sciences de la Vie 319(7):647-651.
- Austin, G.T. 1998a. New subspecies of Nymphalidae (Lepidoptera) from Nevada and Arizona. Pp. 573-586 in: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida. 878 pp.
- Austin, G.T. 1998b. *Callophrys* (Lepidoptera: Lycaenidae) in Nevada, with description of a new subspecies of *C. comstocki*. Pp. 619-628 in: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida, 878 pp.

Austin, G.T. & J.F. Emmel. 1998. New subspecies of butterflies (Lepidoptera) from Nevada and California. Pp. 501-522 *in*: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida, 878 pp.

Austin, G.T., N.M. Haddad, C. Mendez, T.D. Sisk, D.D. Murphy, A.E. Launer & P. R. Ehrlich. 1996. Preliminary annotated checklist of the butterflies of Tikal National Park and vicinity. Tropical Lepidoptera 7(1):21-37.

Austin, G.T. & O.H.H. Mielke. 2000. Hesperiidae of Rondônia, Brazil: *Cephise* Evans (Pyrginae), with descriptions of new species from Mexico and Brazil. Revista Brasileira de Zoologia 17(3):757-788.

Austin, G.T. & D.D. Murphy. 1998. Patterns of phenotypic variation in the *Euphydryas chalcedona* complex (Lepidoptera: Nymphalidae) of the southern intermountain region. Pp. 419-432 *in*: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida, 878 pp.

Austin, G.T. & M.J. Smith. 1998. Revision of the *Thessalia theona* complex (Lepidoptera: Nymphalidae: Melitaeinae). Pp. 359-396 *in*: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida, 878 pp.

Austin, G.T. & A.D. Warren. 2001. Taxonomic notes on some Neotropical Skippers (Lepidoptera: Hesperiidae): *Pyrgus*, *Heliopyrgus*, and *Heliopetes* (Pyrginae). Dugesiana 8(1):1-13.

Austin, G.T. & A.D. Warren. 2002. Taxonomic notes on some Neotropical skippers (Lepidoptera: Hesperiidae): Pyrrhopyginae and Pyrginae. Dugesiana 9(2):15-49.

Back, W. 1991 (1990). Taxonomische untersuchungen innerhalb der Artengruppe um *Euchloe ausonia* (Hübner, 1804) (Lepidoptera, Pieridae). Atalanta 21(3/4):187-206, 318-321.

Bailowitz, R.A. 1988. Systematics of *Ascia* (*Ganya*) (Pieridae) populations in the Sonoran desert. Journal of the Research on the Lepidoptera 26(1-4):73-81.

Bailowitz, R.A. & J.P. Brock. 1991. Butterflies of Southeastern Arizona. Sonoran Arthropod Studies, Inc., Tucson, Arizona. 342 pp.

Bálint, Z. & K. Johnson. 1995. Neotropical polyommatus diversity and affinities. I. Relationships of the higher taxa (Lepidoptera: Lycaenidae). Acta Zoologica Academiae Scientiarum Hungaricae 41(3):211-235.

Bálint, Z. & K. Johnson. 1997. Reformation of the Polyommatus section with a taxonomic and biogeographic overview (Lepidoptera, Lycaenidae, Polyommatini). Neue Entomologische Nachrichten 40:3-68.

Ballmer, G.R. & G.F. Pratt. 1992. *Loranthomitoura*, a new genus of Eumaeini (Lepidoptera: Lycaenidae: Theclinae). *Tropical Lepidoptera* 3(1):37-46.

Basham, B., J. Rathjen, E. Knudson, C. Bordelon, & A. Warren. 2005. *Heliopetes sublinea* (Hesperiidae: Pyrginae), a new record for Texas and the United States. *News of the Lepidopterists' Society* 47(1): 8-10.

Beck, A.F. 1985. Zone 9 Southeast: Tennessee, Mississippi, Alabama, Virginia, North Carolina, South Carolina, Georgia, Florida, Bermuda. Pp. 28-29 in: *Season Summary 1984*. *News of the Lepidopterists' Society* 1985(2):13-33.

Belik, A.G. & D.G. Zamolodchikov. 2002. Notes on systematics of the *Erebia dabanensis* species complex, with special consideration of the *dabanensis-youngi* and *anyuica-occulta* pairs of sibling species (Nymphalidae: Satyrinae). *Nota Lepidopterologica* 25(1):61-79.

Bell, E.L. 1941. On *Lerodea telata* (Herrich-Schäffer) and *tyrtaeus* Ploetz (Lepidoptera: Hesperiidae). *Entomological News* 52(7):183-185.

Bennett, R. & E.C. Knudson. 1976. Additional new butterfly records from Florida. *Journal of the Lepidopterists' Society* 30(3):234-235.

Beutelspacher, C.R. 1986. Estado taxonómico actual del género *Ascia* Scopoli (1777) (Lepidoptera: Pieridae) en México. *Anales del Instituto de Biología, Universidad Nacional Autónoma de México* 57(Ser. Zool. 2):311-322.

Bird, C.D., G.J. Hilchie, N.G. Kondla, E.M. Pike, & F.A.H. Sperling. 1995. Alberta Butterflies. The Provincial Museum of Alberta, Edmonton. 349 pp.

Blum, M. J., E. Beringham & K. Dasmahapatra. 2002. A molecular phylogeny of the neotropical butterfly genus *Anartia* (Lepidoptera: Nymphalidae). *Molecular Phylogenetics and Evolution* 26:46-55.

Bordelon, C.W., Jr. 1991. A note about an important observation. *News of the Lepidopterists' Society* 1991(4):58.

Bordelon, C.W., Jr. 2000. Zone 6 South Central: Oklahoma, Texas, Arkansas, Louisiana. Pp. 33-50 in: *Season Summary 1999*. *News of the Lepidopterists' Society* 42(supplement S1):3-78.

Bordelon, C.W., Jr., & E. Knudson. 2000. New records of Lepidoptera from Texas and the USA, and illustrations of other interesting species. *News of the Lepidopterists' Society* 42(1):3-7, 19.

Bordelon, C.W., Jr., & E. Knudson. 2006. Three new USA butterfly records (Pieridae, Nymphalidae) from the lower Rio Grande Valley of Texas. *News of the Lepidopterists' Society*

48(1): 3-6.

Bowden, S.R. 1972. *Pieris napi* L. (Pieridae) and the superspecies concept. Journal of the Lepidopterists' Society 26(3):170-173.

Boyd, B.M., B.M. Boyd, G.T. Austin & D.D. Murphy. 1999. Hybridization of *Limenitis* in the western Great Basin (Lepidoptera: Nymphalidae). Holarctic Lepidoptera 6(2):37-74.

Brock, J.P. 1998. Life styles of the scaled and beautiful: Nabokov's satyr. American Butterflies 6(3):36-39.

Brower, A.V.Z. 1994. The case of the missing h: *Heliconius charithonia* (L., 1767), not "*Heliconius charitonia* (L., 1767)." Journal of the Lepidopterists' Society 48(2):166-168.

Brower, A.V.Z. 1999. Delimitation of phylogenetic species with DNA sequences: a critique of Davis and Nixon's population aggregation analysis. Systematic Biology 48(1):199-213.

Brower, A.V.Z. 2000. Phylogenetic relationships among the Nymphalidae (Lepidoptera), inferred from partial sequences of the Wingless gene. Proceedings of the Royal Society of London B 267:1201-1211.

Brown, F.M. 1929. A revision of the genus *Phoebis* (Lepidoptera). American Museum Novitates 368:1-22.

Brown, F.M. 1931. A revision of the genus *Aphrissa*. American Museum Novitates 454:1-14.

Brown, F.M. 1955. Studies of Nearctic *Coenonympha tullia* (Rhopalocera, Satyridae). *Coenonympha tullia inornata* Edwards. Bulletin of the American Museum of Natural History 105(4):359-410.

Brown, F.M. 1961. *Coenonympha tullia* on the islands in the St. Lawrence River. The Canadian Entomologist 93(2):107-117.

Brown, F.M. & B. Heineman. 1972. Jamaica and its Butterflies. F.W. Classey Ltd., London. xv + 478 pp., 10 pls.

Brown, J.W. 1983. A new species of *Mitoura* Scudder from southern California (Lepidoptera: Lycaenidae). Journal of the Research on the Lepidoptera 21(4):245-254.

Brown, J.W. 1993. Thorne's Hairstreak, *Mitoura thornei* Brown. Pp. 122-123 in: New, T.R., editor. Conservation biology of Lycaenidae (Butterflies). Occasional paper of the IUCN Species Survival Commission No. 8. Gland, Switzerland. viii + 173 pp.

Brown, J.W., H.G. Real, & D.K. Faulkner. 1992. Butterflies of Baja California. Lepidoptera Research Foundation, Beverly Hills, California. iv + 129 pp., 8 pls.

Brown, K.S., Jr. 1992. Borboletas da Serra do Japi: diversidade, hábitats, recursos alimentares e variação temporal. Pp. 142-187 *in*: Morellato, L.P.C., org. História Natural da Serra do Japi, Ecología e Preservação de uma Área Florestal no Sudeste do Brasil. Universidade Estadual de Campinas, Brazil. 321 pp.

Brussard, P.F., J.F. Baughman, P.D. Murphy, P.R. Ehrlich & J. Wright. 1989. Complex population differentiation in checkerspot butterflies. Canadian Journal of Zoology 38(2):126-143.

Burns, J.M. 1964. Evolution in skipper butterflies of the genus *Erynnis*. University of California Publications in Entomology 37:iv, 1-216.

Burns, J.M. 1974. The polytypic genus *Celotes* (Lepidoptera: Hesperiidae: Pyrginae) from the southwestern United States and northern Mexico. Psyche 81(1):51-69.

Burns, J.M. 1975. Isozymes in evolutionary systematics. Pp. 49-62 *in*: Markert, C.L., editor. Isozymes, IV. Genetics and evolution. Academic Press, Inc., New York. xxi + 965 pp.

Burns, J.M. 1983. Superspecies *Atrytonopsis ovinia* (*A. ovinia* plus *A. edwardsi*) and the nonadaptive nature of interspecific genitalic differences (Lepidoptera: Hesperiidae). Proceedings of the Entomological Society of Washington 85(2):335-358.

Burns, J.M. 1990. *Amblyscirtes*: problems with species, species groups, the limits of the genus, and genus groups beyond- a look at what is wrong with the skipper classification of Evans (Hesperiidae). Journal of the Lepidopterists' Society 44(1):11-27.

Burns, J.M. 1992. Genitalic recasting of *Poanes* and *Paratrytone* (Hesperiidae). Journal of the Lepidopterists' Society 46(1):1-23.

Burns, J.M. 1994a. Split skippers: Mexican genus *Poanopsis* goes in the *origenes* group- and *Yvretta* forms the *rhesus* group- of *Polites* (Hesperiidae). Journal of the Lepidopterists' Society 48(1):24-45.

Burns, J.M. 1994b. Genitalia at the generic level: *Atrytone* restricted, *Anatrytone* resurrected, new genus *Quasimellana*- and yes! we have no *Mellanias* (Hesperiidae). Journal of the Lepidopterists' Society 48(4):273-337.

Burns, J.M. 1996. Genitalia and the proper genus: *Codatractus* gets *mysie* and *uvydixa*- in a compact *cyda* group- as well as a *hysterectomy*, while *Cephise* gets part of *Polythrix* (Hesperiidae: Pyrginae). Journal of the Lepidopterists' Society 50(3):173-216.

Burns, J.M. 2001. *Pyrgus communis* and *Pyrgus albescens* (Hesperiidae) are separate transcontinental species with variable but diagnostic valves. Journal of the Lepidopterists' Society 54(2):52-71.

Burns J.M & D.H. Janzen. 2001. Biodiversity of Pyrrhopygine skipper butterflies (Hesperiidae) in the Area de Conservación de Guanacaste, Costa Rica. Journal of the Lepidopterists' Society 55(1):15-43.

Calhoun, J.V. 1997. Updated list of the butterflies and skippers of Florida (Lepidoptera: Papilionoidea and Hesperioidea). Holarctic Lepidoptera 4(2):39-50.

Callaghan, C.J. & G. Lamas. (2004). Riodinidae. Pages 141-170 *in*: Lamas, G., editor. Checklist: Part 4A. Papilionoidea – Hesperioidea *in*: Heppner, J.B., editor. Atlas of Neotropical Lepidoptera. Scientific Publishers, Gainesville, Florida.

Campbell, D., Brower, A.V.Z. & N.E. Pierce. 2000. Molecular evolution of the Wingless gene and its implications for the phylogenetic placement of the butterfly family Riodinidae (Lepidoptera: Papilionoidea). Molecular Biology and Evolution 17:684-696.

Cary, S.J. & R. Holland. 1992. New Mexico butterflies: checklist, distribution and conservation. Journal of the Research on the Lepidoptera 31(1-2):57-82.

Cassie, B., J. Glassberg, A. Swengel & G. Tudor. 2001. Checklist & English Names of North American Butterflies. North American Butterfly Association, Morristown, New Jersey. 59 pp.

Catling, P.M. & J.V. Calhoun. 1997. Genus *Megisto* in Florida and the taxonomic status of *Megisto viola* (Lepidoptera: Nymphalidae: Satyrinae). Holarctic Lepidoptera 4(1):27-33.

Chermock, R.L. 1950. A generic revision of the Limenitini of the World. American Midland Naturalist 43(3):513-569.

Chew, F.S. & W.B. Watt. 2006. The green-veined white (*Pieris napi* L.), its pierine relatives, and the systematics dilemmas of divergent character sets (Lepidoptera, Pieridae). Biological Journal of the Linnean Society, London. In press.

Chiba, H. & H. Tsukiyama. 1996. A review of the genus *Ochlodes* Scudder, 1872, with special reference to Eurasian species (Lepidoptera: Hesperiidae). Butterflies 14(1):3-16.

Chuah, H.H. & D.S. Cushing. 1995. *Eurema albula* (Pieridae) and *Anthanassa argentea* (Nymphalidae): new records for the United States (Lepidoptera: Papilionoidea). Tropical Lepidoptera 6(1):43-44.

Clench, H.K. 1961. Tribe Theclini, Hairstreaks. Pp. 177-220 *in*: Ehrlich, P.R. & A.E. Ehrlich. How to Know the Butterflies. Wm. C. Brown Co., Dubuque, Iowa. [v] + 262 pp.

Clench, H.K. 1972a. A review of the genus *Lasaia* (Riodinidae). Journal of Research on the Lepidoptera 10(2):149-180.

Clench, H.K. 1972b. *Celastrina ebenina*, a new species of Lycaenidae (Lepidoptera) from the eastern United States. Annals of Carnegie Museum 44(4):33-44.

Clench, H.K. 1977. A list of the butterflies of Andros, Bahamas (Lepidoptera). Annals of Carnegie Museum 46(12):173-194.

Clench, H.K. 1981. New *Callophrys* (Lycaenidae) from North and Middle America. Bulletin of the Allyn Museum 64:1-32.

Comstock, J.A. 1927. Butterflies of California. A Popular Guide to a Knowledge of the Butterflies of California Embracing all of the 477 Species and Varieties at Present Recorded from the State. Author, Los Angeles, California. 334 pp., 63 pls.

Comstock, W.P. 1943. Notes on the subgenus *Glutophrissa*, genus *Appias* (Lepidoptera, Pieridae). American Museum Novitates 1238:1-6.

Comstock, W.P. 1944. The insects of Porto Rico and the Virgin Islands. Rhopalocera or butterflies, 12(4):421-622. In: Scientific Study of Porto Rico and the Virgin Islands. New York Academy of Sciences.

Comstock, W.P. & E.I. Huntington. 1943. Lycaenidae of the Antilles (Lepidoptera, Rhopalocera). Annals of the New York Academy of Sciences 45(2):49-130.

Cowan, C.F. 1970. Annotationes Rhopalocerologicae. Clunbury Press, Berkamsted, England. 70 pp.

Crosson du Cormier, A. 1977. Les *Boloria* du domaine Arctique nécessite d'une nouvelle division spécifique. Alexanor 10(1):35-47.

D'Abrera, B. 1994. Butterflies of the Neotropical Region. Part VI. Riodinidae. Hill House, Victoria, Australia. viii + pp. 880-1096.

d'Almeida, R.F. 1939a. Revisão do gênero *Appias* (subgên. *Glutophrissa* Butl.) (Lepidoptera). Boletim Biológico (n.s.) 4(1):50-66.

d'Almeida, R.F. 1939b. Revisão do gênero *Aphrissa* Butl. (Lepid., Pieridae). Boletim Biológico (n.s.) 4(3):423-443.

d'Almeida, R.F. 1940. Revisão do gênero *Phoebis* Hübn. (Lepidopt. Pieridae). Arquivos de Zoologia do Estado de São Paulo 1(3):67-152.

Dasmahapatra, K. K., M. J. Blum, A. Aiello, S. Hackwell, N. Davies, E. P. Bermingham & J. Mallet. 2002. Inferences from a rapidly moving hybrid zone. Evolution 56(4):741-753.

Dauphin, J., D. Dauphin, E. Knudson, & C. Bordelon. 2005. Three new butterflies new to the

United States and South Texas. News of the Lepidopterists' Society 47(2): 43-46.

Davenport, D. 1941. The butterflies of the satyrid genus *Coenonympha*. Bulletin of the Museum of Comparative Zoology 87(4):215-349.

Davis, K., M. Stangeland, & A.D. Warren. 2005. *Mylon pelopidas*: a new record for Arizona and the United States. News of the Lepidopterists' Society 47(4): 103-104, 106, 110.

de Jong, R. 1972. Systematics and geographic history of the genus *Pyrgus* in the Palaearctic region (Lepidoptera, Hesperiidae). Tijdschrift voor Entomologie 115(1):1-121.

de Jong, R. 1978. Monograph of the genus *Spialia* Swinhoe (Lepidoptera, Hesperiidae). Tijdschrift voor Entomologie 121(3):23-146.

de Jong, R., R.I. Vane-Wright & P.R. Ackery. 1996. The higher classification of butterflies (Lepidoptera): problems and prospects. Entomologica Scandanavica 27(1):65-101.

de la Maza, R.E., J.E. de la Maza & A.L. White. 1989. La fauna de mariposas de México. Revista de la Sociedad Mexicana de Lepidopterología 12(2):39-98.

de la Maza, R.E. & R.D. Turrent. 1985. Mexican Lepidoptera. Eurytelinae I. Publicaciones Especiales de la Sociedad Mexicana de Lepidopterología (Mexico City). 44 pp., [30]pls.

de Lesse, H. 1952. Note sur les genres *Precis* Hb. et *Junonia* Hb. (Lep. Nymphalidae). Bulletin, Société Entomologique du France 57(5):74-77.

DeVries, P.J. 1987. The Butterflies of Costa Rica and their Natural History. Papilionidae, Pieridae, Nymphalidae. Princeton University Press, Princeton, New Jersey. xxii + 327 pp., 50 pls.

DeVries, P.J. 1997. The Butterflies of Costa Rica and their Natural History. Volume II: Riodinidae. Princeton University Press, Princeton, New Jersey. xxv + 288 pp., 25 pls.

Devyatkin, A.L. 1990. [On two Siberian species of *Pyrgus*] (Lepidoptera, Hesperiidae). Zoologicheskii Zhurnal 69(10):141-145.

Dimock, T.E. 1973. Three natural hybrids of *Vanessa atalanta rubria* x *Cynthia annabella* (Nymphalidae). Journal of the Lepidopterists' Society 27(4):274-278.

dos Passos, C.F. 1958. The satyrid butterflies of northwestern North America. Proceedings of the Tenth International Congress of Entomology 1:673-682.

dos Passos, C.F. 1960. Taxonomic notes on some Nearctic Rhopalocera. 1. Hesperioidea. Journal of the Lepidopterists' Society 14(1):24-36.

dos Passos, C.F. 1961. Taxonomic notes on some Nearctic Rhopalocera. 2. Papilionoidea. Journal of the Lepidopterists' Society 15(4):209-225.

dos Passos, C.F. 1964. A synonymic list of the Nearctic Rhopalocera. Memoirs of the Lepidopterists' Society 1:v, 1-145.

dos Passos, C.F. & L.P. Grey. 1945. A genitalic survey of the Argynninae (Lepidoptera, Nymphalidae). American Museum Novitates 1296:1-29.

Dubatolov, V.V. 1992. [New subspecies of nymphalid and satyrid butterflies (Lepidoptera, Rhopalocera) from Yakutia.] Vestnik Zoologii 1992(6):40-45.

Dubatolov, V.V. & O.E. Kosterin. 1994. A new subspecies of the *Euchloe ausonia* Hübner species group from the highlands of the Dzhungerian Alatau (East Kazakhstan). Atalanta 25(3/4):513-520.

Durden, C.J. 1982. The butterfly fauna of Barton Creek Canyon on the Balcones Fault Zone, Austin, Texas, and a regional list. Journal of the Lepidopterists' Society 36(1):1-17.

Eitschberger, U. 1981. Die nordamerikanischen Arten aus der *Pieris napi-bryoniae*-Gruppe (Lep. Pieridae). Atalanta 11(5):366-371.

Eitschberger, U. 1983. Systematische Untersuchungen am *Pieris napi-bryoniae*-Komplex (s.l.) (Lepidoptera, Pieridae). Herbipoliana 1(1):i-xxii, 1-504; 1(2):1-601.

Eitschberger, U. 1993. Die Struktur der Eihüllen einiger *Papilio*-Arten im Vergleich unter dem REM/SEM. Atalanta 24(1/4):15-32.

Eliot, J.N. 1973. The higher classification of the Lycaenidae (Lepidoptera): a tentative arrangement. Bulletin of the British Museum of Natural History (Entomology) 28(6):1-505.

Eliot, J.N. 1990. Notes on the genus *Curetis* Hübner (Lepidoptera, Lycaenidae). Transactions of the Lepidopterological Society of Japan 41: 201-205.

Eliot, J.N. & A. Kawazoé. 1983. Blue Butterflies of the *Lycaenopsis* Group. British Museum (Natural History), London. 309 pp.

Emmel, J.F. & T.C. Emmel. 1998. A new *Agriades* (Lepidoptera: Lycaenidae) from the Sierra Nevada and Trinity Alps of California, and the biology and geographic variation of *Agriades podarce* in California. Pp. 287-302 in: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida. 878 pp.

Emmel, J.F., T.C. Emmel, & S.O. Mattoon. 1998a. The types of butterflies named by Jean Alphonse Boisduval: designation of lectotypes and a neotype, and fixation of type localities. Pp. 3-76 in: Emmel, T.C., editor. 1998. Systematics of Western North American Butterflies.

Mariposa Press, Gainesville, Florida. 878 pp.

Emmel, J.F., T.C. Emmel, & S.O. Mattoon. 1998b. New subspecies of Pieridae (Lepidoptera) from California, Nevada, and Baja California. Pp. 127-138 in: Emmel, T.C., editor. 1998. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida. 878 pp.

Emmel, T.C., editor. 1998. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida. 878 pp.

Emmel, T.C. & G.T. Austin. 1998. What is *Argynnис carolae*? The cytotaxonomy and systematic position of a *Speyeria* relict from Nevada (Lepidoptera: Nymphalidae). Pp. 443-450 in: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida, 878 pp.

Emmel, T.C. & J.F. Emmel. 1973. The Butterflies of Southern California. Natural History Museum of Los Angeles County, Science Series 26. xi + 148 pp.

Evans, W.H. 1951. A Catalogue of the Hesperiidae Indicating the Classification and Nomenclature Adopted in the British Museum (Natural History). Part I. Pyrrhopyginae. British Museum of Natural History, London. 92 pp., pls. 1-9.

Evans, W.H. 1952. A Catalogue of the Hesperiidae Indicating the Classification and Nomenclature Adopted in the British Museum (Natural History). Part II. Pyrginae. Section I. British Museum of Natural History, London. 178 pp., pls. 10-25.

Evans, W.H. 1953. A Catalogue of the Hesperiidae Indicating the Classification and Nomenclature Adopted in the British Museum (Natural History). Part III. Pyrginae. Section II. British Museum of Natural History, London. 246 pp., pls. 26-53.

Evans, W.H. 1955. A Catalogue of the Hesperiidae Indicating the Classification and Nomenclature Adopted in the British Museum (Natural History). Part IV. Hesperiinae and Megathyminae. British Museum of Natural History, London. 499 pp., pls. 54-88.

Faynel & Bálint, Z. 2004. Bulletin de la Societe Entomologique France 109: 263-286.

Ferris, C.D. 1977. Taxonomic revision of the species *dorcas* Kirby and *helloides* Boisduval in the genus *Epidemia* Scudder. Bulletin of the Allyn Museum 45:1-42.

Ferris, C.D. 1981. A new subspecies of *Colias alexandra* W.H. Edwards and notes on *Colias hecla* Lefebvre (Pieridae: Coliadinae). Bulletin of the Allyn Museum 63:1-12.

Ferris, C.D. 1983. *Speyeria atlantis* phenotypes in the southern Rocky Mountains (Lepidoptera: Nymphalidae: Argynninae). Journal of Research on the Lepidoptera 22(2):101-114.

Ferris, C.D. 1988. Revision of several North American Leguminosae-feeding *Colias* species, with description of a new subspecies (Pieridae: Coliadinae). Bulletin of the Allyn Museum 116:1-28.

Ferris, C.D. 1989a. A new species of *Colias* from Utah (Pieridae: Coliadinae). Bulletin of the Allyn Museum 128:1-11.

Ferris, C.D. 1989b (1988). *Euphydryas anicia* and *E. chalcedona* in Idaho (Lepidoptera: Nymphalidae). Journal of Research on the Lepidoptera 26(1-4):109-115.

Ferris, C.D., editor. 1989c. Supplement to: A catalogue/checklist of the butterflies of America north of Mexico. Memoirs of the Lepidopterists' Society 3:viii, 1-103.

Ferris, C.D. 1991. Appearance of the "heathii" aberration and genitalic variation in a *Mitoura* population from Oregon (Lycaenidae: Theclinae). Journal of Research on the Lepidoptera 30(1-2):113-120.

Ferris, C.D. 1993. Reassessment of the *Colias alexandra* group, the legume-feeding species, and preliminary cladistic analysis of the North American *Colias* (Pieridae: Coliadinae). Bulletin of the Allyn Museum 138:1-91.

Field, W.D. 1940 (1938). A manual of the butterflies and skippers of Kansas. Bulletin of the University of Kansas. Biological Series (Department of Entomology, No. 12). 39(10):1-328.

Field, W.D. 1971. Butterflies of the genus *Vanessa* and the resurrected genera *Bassaris* and *Cynthia* (Lepidoptera- Nymphalidae). Smithsonian Contributions to Zoology 84:1-105.

Fisher, M.S. 1977. The taxonomy and identity of *Papilio nitra* W.H. Edwards in Colorado. Bulletin of the Allyn Museum 47:1-8.

Forbes, W.T.M. 1960. Lepidoptera of New York and neighboring states, part IV. Agaristidae through Nymphalidae including butterflies. Cornell University Agricultural Experiment Station Memoir 371:5-188.

Forster, W. 1964. Beiträge zur Kenntnis der Insektenfauna Boliviens XIX. Lepidoptera III, Satyridae. Veröffentlichungen der Zoologischen Staatssammlung München 8:51-188.

Freeman, H.A. 1946. Two new species of skippers from North and Central America. Entomological News 57(8):185-187.

Freeman, H.A. 1951. Notes on the genus *Yvretta* Hemming with a new record for the United States. Field and Laboratory 19(1):45-46.

Freeman, H.A. 1959. A revision of the genera of the Megathyminidae, with descriptions of three new genera. The Lepidopterists' News 12(3/4):81-92.

- Freeman, H.A. 1960. Butterfly collecting in Texas and New Mexico. *Journal of the Lepidopterists' Society* 13(2):89-93.
- Freeman, H.A. 1964a. A new species of *Apodemia* from Texas (Riodinidae). *Journal of the Lepidopterists' Society* 18(2):75-77.
- Freeman, H.A. 1964b. Four new species of *Agathymus* from Texas (Megathyminidae). *Journal of the Lepidopterists' Society* 18(3):171-185.
- Freeman, H.A. 1967. *Polythrix octomaculata*, not *procerus*, in Texas (Hesperiidae). *Journal of the Lepidopterists' Society* 21(4):278.
- Freeman, H.A. 1969. Records, new species, and a new genus of Hesperiidae from Mexico. *Journal of the Lepidopterists' Society* 23(supplement 2):1-62.
- Freeman, H.A. 1970. A new species of *Piruna* from Texas (Hesperiidae). *Journal of the Lepidopterists' Society* 24(4):247-249.
- Freeman, H.A. 1975. A new species of *Euphyes* from Texas (Hesperiidae). *Journal of the Lepidopterists' Society* 29(4):227-229.
- Freeman, H.A. 1977. *Staphylus azteca*, new record for the United States (Hesperiidae). *Journal of the Lepidopterists' Society* 31(1):62.
- Freeman, H.A. 1993. Notes on *Amblyscirtes* Scudder, with the description of two new subspecies (Insecta: Lepidoptera: Hesperiidae: Hesperiinae). *Annals of Carnegie Museum* 62(4):341-350.
- Freeman, H.A. & A.D. Warren. (in preparation). Revision of the genus *Piruna* Evans.
- Freitas, A.V.L. 1999. Nymphalidae (Lepidoptera), filogenia com base en caracteres de imaturos, com experimentos de troca de plantas hospedeiras. Universidade Estadual de Campinas, Campinas, São Paulo, Brazil. xii + 170 pp.
- Freitas, A.V.L. & K.S. Brown, Jr. (in preparation). Phylogeny of the Nymphalidae (Lepidoptera).
- Friedlander, T. 1984. Another look at snout butterflies (Libytheidae: *Libytheana*). *Journal of the Lepidopterists' Society* 38(2):139-141.
- Friedlander, T. 1987. Taxonomy, phylogeny, and biogeography of *Asterocampa* Röber 1916 (Lepidoptera, Nymphalidae, Apaturinae). *Journal of Research on the Lepidoptera* 25(4):1-337.

Gall, L.F. & F.A.H. Sperling. 1980. A new high altitude species of *Boloria* from southwestern Colorado (Nymphalidae), with a discussion of phenetics and hierarchical decisions. Journal of the Lepidopterists' Society. 34(2):230-252.

Gatrell, R.R. 1998. The rediscovery, taxonomy, and biology of *Chlosyne gorgone gorgone* and *Chlosyne ismeria* (Nymphalidae) in Burke County, Georgia. The Taxonomic Report 1(2):1-9.

Gatrell, R.R. 1999. Hübner's *helicta*: the forgotten *Neonympha*. The Taxonomic Report 1(8):1-8.

Gatrell, R.R. 2000. *Chlosyne ismeria ismeria*. The International Lepidoptera Survey Newsletter 2(3):1.

Gatrell, R.R. 2004. Description of a multilevel cryptic new species of *Phyciodes* (Nymphalidae: Melitaeinae) from the southern Appalachian mountains. The Taxonomic Report 4(8): 1-19.

Geiger, H. & A.M. Shapiro. 1986. Electrophoretic evidence for speciation within the nominal species *Anthocharis sara* Lucas (Pieridae). Journal of Research on the Lepidoptera 25(1):15-24.

Geiger, H. & A.M. Shapiro. 1992. Genetics, systematics, and evolution of holarctic *Pieris napi* group populations (Lepidoptera: Pieridae). Zeitschrift für Zoologisches Systematik und Evolutionforschung 30(2):100-122.

Glassberg, J. 1992. *Aglais urticae* (Nymphalidae): a nascent population in North America. Journal of the Lepidopterists' Society 46(4):302-304.

Glassberg, J. 2004. Go get set on your marks: "Green longtails." American Butterflies 12(1): 28-38.

Gochfield, M. & J. Burger. 1997. Butterflies of New Jersey. Rutgers University Press, New Brunswick, New Jersey. 327 pp.

Goodpasture, C. 1973. Biology and systematics of the *Plebejus (Icaricia) acmon* group (Lepidoptera: Lycaenidae). I. Review of the group. Journal of the Kansas Entomological Society 46(4):468-485.

Goodpasture, C. 1974. Foodplant specificity in the *Plebejus (Icaricia) acmon* group (Lycaenidae). Journal of the Lepidopterists' Society 28(1):53-63.

Gorbunov, P.Y. 2001. The Butterflies of Russia: Classification, Genitalia, Keys for Identification (Lepidoptera: Hesperioida and Papilioidea). Russian Academy of Sciences, Institute of Plant and Animal Ecology, "Thesis", Ekaterinburg, Russia. 320 pp.

Grey, L.P. 1957. Review. Warren's argynnid classification (Nymphalidae). The Lepidopterists' News 11(4-5):171-176.

Grey, L.P. 1989. Sundry argynnine concepts revisited (Nymphalidae). Journal of the Lepidopterists' Society 43(1):1-10.

Grishin, N.V. 2005. A new banner for the United States: *Temenis laothoe* (Nymphalidae: Biblidinae). New of the Lepidopterists' Society 47 (1): 3, 10.

Guppy, C.S. & J.H. Shepard. 2001. Butterflies of British Columbia. University of British Columbia Press, Vancouver. 414 pp.

Hafernik, J.E. 1982. Phenetics and ecology of hybridization in buckeye butterflies (Lepidoptera: Nymphalidae). University of California Publications in Entomology 96:1-109.

Hagen, R.H., R.C. Lederhouse, J.L. Bossart, & J.M. Scriber. 1991. *Papilio canadensis* and *P. glaucus* (Papilionidae) are distinct species. Journal of the Lepidopterists' Society 45(4):245-258.

Hammond, P.C. & D.V. McCorkle. 2000. A new species of *Philotiella* from the Oregon Cascade Range (Lepidoptera: Lycaenidae). Holarctic Lepidoptera 6(2):77-82.

Handfield, L. 1999. Le Guide des Papillons du Québec. Volume I (version scientifique). Broquet, Boucherville, Canada. 982 pp., 122 pls.

Hanson, D.J., E. Knudson, & C. Bordelon. 2003. *Phocides belus* Godman and Salvin (Hesperiidae), new to US and Texas with a review of similar species of the USA and northern Mexico.

Harvey, D.J. 1991. Higher classification of the Nymphalidae. Pp. 255-273 in: Nijhout, H.F. The Development and Evolution of Butterfly Wing Patterns. Smithsonian Institution Press, Washington, D.C. xvi + 297 pp.

Hassler, M. & H. Feil. 2002. A Siberian butterfly recorded for the first time on the American continent: *Oeneis nanna kluanensis* ssp. n. (Lepidoptera: Nymphalidae, Satyrinae). Nachrichten Entomologischer Verein Apollo, N.F. 22(4):197-205.

Heitzman, J.R. 1973. A new species of *Papilio* from the eastern United States. Journal of Research on the Lepidoptera 12(1):1-10.

Heitzman, J.R. & R.L. Heitzman. 1973. New Butterfly Records for the United States (Hesperiidae & Libytheidae). Journal of Research on the Lepidoptera 10(4):284-286.

Hemming, F. 1934. The generic names of the Holarctic butterflies. Volume I: 1758-1863. The British Museum (Natural History), London. viii + 184 pp.

Hemming, F. 1967. The generic names of the butterflies and their type-species (Lepidoptera: Rhopalocera). Bulletin of the British Museum of Natural History (Entomology) 9:1-509.

Hesselbarth, G., Van Oorschot, H. & S. Wagener. 1995. Die tagfalter der Türkei. 3 volumes. S. Wagener, Bocholt, Germany.

Higgins, L.G. 1960. A revision of the melitaeine genus *Chlosyne* and allied species (Lepidoptera: Nymphalidae). Transactions of the Royal Entomological Society of London 112(14):381-475.

Higgins, L.G. 1978. A revision of the genus *Euphydryas* Scudder (Lepidoptera: Nymphalidae). Entomologist's Gazette 29(3):109-115.

Higgins, L.G. 1981. A revision of *Phyciodes* Hübner and related genera, with a review of the classification of the Melitaeinae (Lepidoptera: Nymphalidae). Bulletin of the British Museum of Natural History (Entomology) 43(3):77-243.

Higgins, L.G. & N.D. Riley. 1970. A Field Guide to the Butterflies of Britain and Europe. Collins, London. 380 pp., 60 pls.

Hilchie, G.J. 1990. Classification, relationships, life history, and evolution of *Erebia magdalena* Strecker (Lepidoptera, Satyridae). Quaestiones Entomologicae 26(4):665-693.

Hinchliff, J. 1994. An Atlas of Oregon Butterflies. Oregon State University Bookstore, Corvallis, Oregon. v + 176 pp.

Hinchliff, J. 1996. An Atlas of Washington Butterflies. Oregon State University Bookstore, Corvallis, Oregon. vi + 162 pp.

Holland, R. & G.S. Forbes. 1981. Rediscovery of *Apodemia phyciodoides* (Riodinidae). Journal of the Lepidopterists' Society 35(3):226-232.

ICZN (International Commission on Zoological Nomenclature). 1999. International Code of Zoological Nomenclature, Fourth Edition. The International Trust for Zoological Nomenclature, London. xxix + 306 pp.

ICZN (International Commission on Zoological Nomenclature). 2006. Opinion 2144 (Case 3282). *Thecla azia* Hewitson, 1873 (Insecta, Lepidoptera): specific name conserved. Bulletin of Zoological Nomenclature 63(1): 76-77.

Janzen, D.H., M.J. Sharkey, & J.M. Burns. 1998. Parasitization biology of a new species of Braconidae (Hymenoptera) feeding on larvae of Costa Rican dry forest skippers. Tropical Lepidoptera 9 (supplement 2):33-41.

Jenkins, D.W. 1983. Neotropical Nymphalidae I. Revision of *Hamadryas*. Bulletin of the Allyn Museum 81:1-146.

- Johnson, K. 1976. Three new Nearctic species of *Callophrys* (*Mitoura*), with a diagnostis [sic] of all Nearctic consubgenera (Lepidoptera: Lycaenidae). Bulletin of the Allyn Museum 38:1-30.
- Johnson, K. 1992. The Palaearctic “elfin” butterflies (Lycaenidae, Theclinae). Neue Entomologische Nachrichten 29(1):1-141.
- Johnson, K., G.T. Austin, J.F. Le Crom & J.A. Salazar. 1997. The Strephonina, a new infratribe of the Eumaeini with description of fourteen new genera (Lycaenidae). Revista de Theclinae Colombianos 1(1):[ii], 1-57, 2 pls.
- Johnson, K. & Z. Bálint. 1995. Distinction of *Pseudochrysops*, *Cyclargus*, *Echinargus* and *Hemiargus* in the Neotropical Polyommatus (Lepidoptera, Lycaenidae). Reports of the Museum of Natural History, University of Wisconsin, Stevens Point 54:1-14.
- Johnson, K. & G. Balogh. 1977. Studies in the Lycaeninae. 2. Taxonomy and evolution of the Nearctic *Lycaena rubidus* complex, with description of a new species. Bulletin of the Allyn Museum 43:1-62.
- Johnson, K. & J.F. Le Crom. 1997. Clench’s *Cyanophrys* “amyntor Group” in Colombia with descriptions of several outlying endemics from elsewhere in South America. Revista de Theclinae Colombianos 1(3):[ii], 1-28.
- Karsholt, O. & J. Razowski. 1996. The Lepidoptera of Europe. A Distributional Checklist. Apollo Books, Stenstrup, Germany. 380 pp.
- Kawahara, A.Y. 2001. Systematics of the snout butterflies (Lepidoptera: Nymphalidae: Libytheinae) with an emphasis on morphology. Entomology Department Honors Thesis, Cornell University, Ithaca, New York. 161 pp.
- Kendall, R.O. 1974a. Confirmation of Rhopalocera (Pieridae, Nymphalidae) previously recorded for Texas and the United States. Journal of the Lepidopterists' Society 28(3):249-252.
- Kendall, R.O. & W.W. McGuire. 1984. Some new and rare records of Lepidoptera found in Texas. Bulletin of the Allyn Museum 86:1-50.
- Kimball, C.P. 1965. Arthropods of Florida and neighboring land areas. Florida Department of Agriculture, Gainesville. v + 363 pp., 26 pls.
- Klassen, P., A.R. Westwood, W.B. Preston & W.B. McKillop. 1989. The Butterflies of Manitoba. Manitoba Museum of Man and Nature, Winnipeg. vi + 290 pp.
- Klots, A.B. 1933. A generic revision of the Pieridae (Lepidoptera). Together with a study of the male genitalia. Entomologica Americana (n.s.) 12(3):139-204, (4):205-242; pls. 5-13.

Klots, A.B. 1936. The interrelationships of the species of the genus *Lycaena* Fabricius (Lepidoptera, Lycaenidae). Bulletin of the Brooklyn Entomological Society 31(2):154-171.

Klots, A.B. 1951. A Field Guide to Eastern Butterflies. Houghton Mifflin Co., Boston, Massachusetts. xvi + 349 pp.

Knudson, E. 2002. The lower Rio Grande Valley revisited. Southern Lepidopterists' News 24(4): 79-83, insert A.

Knudson, E., C. Bordelon, and A. Warren. 2004. *Antigonus erosus* Hübner (Hesperiidae, Pyrginae), a new US record from South Texas. News of the Lepidopterists' Society 46(4): 111.113.

Koçak, A.Ö. 1983. On the nomenclature of *Papilio thersamon* Esper, 1784 from the west Palaearctic region (Lycaenidae, Lepidoptera). Priamus 3(1):3-5.

Kondla, N.G. 1999. Pend-d Oreille Butterfly Survey. Living Landscapes Program Report for Royal BC Museum and Columbia Basin Trust. Victoria, British Columbia. 38 pp. (<http://www.livingbasin.com/butterfly/>)

Kondla, N.G. & J.P. Pelham. 1995. First record of the butterfly *Euchloe naina* (Lepidoptera, Pieridae) from North America. Canadian Field-Naturalist 109(2):259.

Kons, H.L. 2000. Phylogenetic studies of the Melitaeini (Lepidoptera, Nymphalidae) and a revision of the genus *Chlosyne* Butler. Ph.D. dissertation, University of Florida, Gainesville. x + 798 pp.

Kosterin, O.E. 2000. Description of a new subspecies of *Boloria pales* ([Denis & Schiffermüller], 1775) sensu Warren (1944) (Lepidoptera, Nymphalidae) from Altai. Far East Entomologist 86(1):1-10.

Krizek, G.O. 1998. *Hemiargus ammon* discovered in the USA at Big Pine Key, Florida (Lepidoptera: Lycaenidae). Holarctic Lepidoptera 5(2):45.

Kudrna, O. & H. Geiger. 1985. A critical review of "Systematische Untersuchungen am *Pieris napi-bryoniae*-Komplex (s.l.)" (Lepidoptera: Pieridae), by Ulf Eitschberger. Journal of Research on the Lepidoptera 24(1):47-60.

Lafontaine, J.D. & D.M. Wood. 1997. Butterflies and moths (Lepidoptera) of the Yukon. Pp. 723-785 in: Danks, H.V. & J.A. Downes, editors. Insects of the Yukon. Biological Survey of Canada, Ottawa. 1034 pp.

Lamas, G. 1981. Notes on Peruvian butterflies (Lepidoptera). VI. Twelve new Pieridae. Revista de Ciencias 73(1):44-53.

Lamas, G., editor. 2004. Checklist: Part 4A. Papilionoidea – Hesperioidea *in:* Heppner, J.B., editor. Atlas of Neotropical Lepidoptera. Scientific Publishers, Gainesville, Florida. xxxvi+439 pp.

Lane, C.P. & S.J. Weller. 1994. A review of *Lycaeides* Hübner and karner blue butterfly taxonomy. Pp. 5-21 *in:* Andow, D.A., R.J. Baker & C.P. Lane, editors. Karner Blue Butterfly, a Symbol of a Vanishing Landscape. Miscellaneous Publication 84-1994, Minnesota Agricultural Experiment Station, University of Minnesota, St. Paul. [vi] + 222 pp.

Larsen, T.B. 1991. The Butterflies of Kenya and their Natural History. Oxford University Press, Oxford. xxii + 490 pp., 64 pls.

Layberry, R.A., P.W. Hall, & J.D. Lafontaine. 1998. The Butterflies of Canada. University of Toronto Press, Toronto. 280 pp., 32 pls.

Lindsey, A.W. 1928. *Pyrgus centaureae* and *freija* (Lep.: Hesperiidae). Entomological News 39:239-242.

Lindsey, A.W., E.L. Bell & R.C. Williams, Jr. 1931. The Hesperioidea of North America. Denison University Bulletin, Journal of the Scientific Laboratories 26(1):1-142.

Llorente, J.B. 1984. Sinopsis sistemática y biogeográfica de los Dismorphiinae de México con especial referencia al género *Enantia* Hübner (Lepidoptera: Pieridae). Folia Entomológica Mexicana 58(1):1-207.

Llorente, J.B., L.O. Oñate, A.M. Luis & I.F. Vargas. 1997. Papilionidae y Pieridae de México: Distribución Geográfica e Ilustración. Comisión Nacional Para el Conocimiento y Uso de la Biodiversidad (CONABIO) y Facultad de Ciencias, Universidad Nacional Autónoma de México, México, D.F. viii + 227 pp., 28 pls.

Luis, A.M., J.B. Llorente & I.F. Vargas. 1996. Distribución geográfica, estacionalidad y comportamiento de *Diaethria asteria* (Lepidoptera: Nymphalidae). Tropical Lepidoptera 7(1):39-44.

Luis, A.M., J.B. Llorente, I.F. Vargas & A.D. Warren. 2003. Biodiversity and biogeography of Mexican butterflies (Lepidoptera: Papilionoidea and Hesperioidea). Proceedings of the Entomological Society of Washington 105(1):209-224.

Lukhtanov, V.A. 1989. [Review of the Palaearctic satyrid group *Oeneis norna* (Lepidoptera, Satyridae). Communication 1.], Vestnik Zoologii 1989(2):29-36.

Lukhtanov, V.A. & U. Eitschberger. 2000. Illustrated catalogue of the genera *Oeneis* and *Davidina* (Lepidoptera, Satyrinae, Oeneini). Part 11, pp. 1-12, 27 pls. *in:* Bauer, E., editor. Butterflies of the World. Goecke & Evers, Keltern, Germany.

MacNeill, C.D. 1964. The skippers of the genus *Hesperia* in western North America, with special reference to California (Lepidoptera: Hesperiidae). University of California Publications in Entomology 35:iv, 1-130.

MacNeill, C.D. 1970. A new *Pholisora* with notes on *P. alpheus* (Edw.) (Lepidoptera: Hesperiidae). Entomological News 81(7):177-184.

MacNeill, C.D. 1975. Family Hesperiidae. Pp. 423-578 in: Howe, W.H., editor. The Butterflies of North America. Doubleday and Co., Garden City, New York. xiii + 633 pp., 97 pls.

Masters, J.H. 1970. Distributional notes on the genus *Mestra* (Nymphalidae) in North America. Journal of the Lepidopterists' Society 24(3):203-208.

Mattoni, R.H.T. 1989 (1988). The *Euphilotes battoides* complex: recognition of a species and description of a new subspecies. Journal of Research on the Lepidoptera 27(3-4):173-185.

Mayden, R.L. 1997. A hierarchy of species concepts: the denouement in the saga of the species problem. Pp. 381-424 in: Claridge, M.F., H.A. Dawah & M.R. Wilson, editors. Species. The Units of Biodiversity. Chapman & Hall, London.

Mayr, E. 1957a. Species concepts and definitions. Pp. 1-22 in: Mayr, E., editor. The Species Problem. American Association for the Advancement of Science. Washington, D. C.

Mayr, E. 1957b. Difficulties and importance of the biological species concept. Pp. 371-388 in: Mayr, E., editor. The Species Problem. American Association for the Advancement of Science. Washington, D. C.

McAlpine, W.S. 1971. A revision of the butterfly genus *Calephelis* (Riodinidae). Journal of Research on the Lepidoptera 10(1):9-125.

McGuire, W.W. & M.A. Rickard. 1976. New Hesperiidae records for Texas and the United States. Journal of the Lepidopterists' Society 30(1):5-11.

Mello, M.J. 1999. Zone 10 Northeast: Maryland, New York, Pennsylvania, New England States, Maritime Provinces. Pp. 68-72 in: Season Summary 1998. News of the Lepidopterists' Society 41(supplement S1):1-78.

Mielke, O.H.H. 2002. Pyrrhopyginae: géneros novos e revalidados (Lepidoptera, Hesperiidae). Revista Brasileira de Zoologia 19(1):217-228.

Mielke, O.H.H. 2004. Hesperioidea. Pages 25-86 in: Lamas, G., editor. Checklist: Part 4A. Papilioidea – Hesperioidea in: Heppner, J.B., editor. Atlas of Neotropical Lepidoptera. Scientific Publishers, Gainesville, Florida.

Mielke, O.H.H. & M.M. Casagrande. 2002. Notas taxonômicas em Hesperiidae neotropicais,

com descrições de novos taxa (Lepidoptera). Revista Brasileira de Zoologia 19(supl. 1):27-76.

Mielke, O.H.H. & A.D. Warren. 2004. The identity of *Eudamus valeriana* Plötz (Lepidoptera, Hesperiidae, Pyrginae). Revista Brasileira de Zoologia 21(2): 307-308.

Miller, J.S. 1987. Phylogenetic studies in the Papilioninae (Lepidoptera: Papilionidae). Bulletin of the American Museum of Natural History 186(4):367-512.

Miller, L.D. 1968. The higher classification, phylogeny and zoogeography of the Satyridae (Lepidoptera). Memoirs of the American Entomological Society 24:iii, 1-174.

Miller, L.D. 1972. Revision of the Euptychiini (Satyridae). 1. Introduction and *Paramacera* Butler. Bulletin of the Allyn Museum 8:1-18.

Miller, L.D. 1974. Revision of the Euptychiini (Satyridae). 2. *Cyllopsis* R. Felder. Bulletin of the Allyn Museum 20:1-98.

Miller, L.D. & F.M. Brown. 1979. Studies in the Lycaeninae (Lycaenidae). 4. The higher classification of the American coppers. Bulletin of the Allyn Museum 51:1-30.

Miller, L.D. & F.M. Brown. 1981. A catalogue/checklist of the butterflies of America north of Mexico. Memoirs of the Lepidopterists' Society 2:vii, 1-280.

Miller, L.D. & J.Y. Miller. 1980. Nearctic *Agelais* and *Nymphalis* (Lepidoptera, Nymphalidae): Laurasia revisited? The Entomologist 109(2):106-115.

Motono, S. 1993. Libytheidae of the world. Kirihara Shoten, Tokyo. 52 pp.

Nabokov, V. 1945. Notes on neotropical Plebejinae (Lycaenidae, Lepidoptera). Psyche 52(1-2):1-61.

Neck, R.W. 1996. A Field Guide to Butterflies of Texas. Gulf Publishing Co., Houston, Texas. 323 pp., 64 pls.

Neild, A.F.E. 1996. The Butterflies of Venezuela, Part 1: Nymphalidae I (Limenitidinae, Apaturinae, Charaxinae). Meridian Publications, Greenwich, London. 144 pp., 32 pls.

Nice, C.C. & A.M. Shapiro. 1999. Molecular and morphological divergence in the butterfly genus *Lycaeides* (Lepidoptera: Lycaenidae) in North America: evidence of recent speciation. Journal of Evolutionary Biology 12:936-950.

Nice, C.C. & A.M. Shapiro. 2001. Population genetic evidence of restricted gene flow between host races in the butterfly genus *Mitoura* (Lepidoptera: Lycaenidae). Annals of the Entomological Society of America 94(2):257-267.

Niculescu, E.V. 1986. Familia Nymphalidae (Lep). Phylogenie et Classification. Bulletin de la Société Entomologique de Mulhouse 1986:1-5.

Nielsen, M.C. 1999. Michigan Butterflies and Skippers. Michigan State University Extension, East Lansing. 248 pp.

Nixon, K.C. & Q.D. Wheeler. 1990. An amplification of the phylogenetic species concept. Cladistics 6(3):211-223.

Nylin, S., K. Nyblom, F. Ronquist, N. Janz, J. Belicek & M. Källersjö. 2001. Phylogeny of *Polygonia*, *Nymphalis*, and related butterflies (Lepidoptera: Nymphalidae): a total-evidence analysis. Zoological Journal of the Linnean Society 137(4):441-468.

Oehlenschlager, R. & R. Huber. 2002. The butterflies of Wadena County, Minnesota. Scientific Publications of the Science Museum of Minnesota, New Series 8(1):iv, 1-26.

Okano, K. 1987. The check list of Libytheid butterflies of the world, with some bibliographical observations. Tokurana (Acta Rhopalocera) 13(1):1-12.

Opler, P.A. 1965. Studies on the Nearctic genus *Euchloe* (Pieridae: Lepidoptera). Masters Thesis, San Jose State University, San Jose, California [California State University at San Jose] 172 pp.

Opler, P.A. 1967a. (1966) Studies on the Nearctic *Euchloe*. Part 3. Complete synonymous treatment. Journal of Research on the Lepidoptera 5(3):185-190.

Opler, P.A. 1967b. (1966) Studies on the Nearctic *Euchloe*. Part 4. Type data and type locality restrictions. Journal of Research on the Lepidoptera 5(3):190-195.

Opler, P.A. 1970. Studies on Nearctic *Euchloe*. Part 5. Distribution. Journal of Research on the Lepidoptera 7(2):65-86.

Opler, P.A. 1971. Studies on Nearctic *Euchloe*. Part 6. Systematics of adults. Journal of Research on the Lepidoptera 8(4):153-168.

Opler, P.A. 1974. Studies on Nearctic *Euchloe*. Part 7. Comparative life histories, hosts and the morphology of the immature stages. Journal of Research on the Lepidoptera 13(1):1-120.

Opler, P.A. 1987. A new *Euchloe* (Pieridae) from northwestern Mexico. Journal of the Lepidopterists' Society 40(3):188-190.

Opler, P.A. 1992. A Field Guide to Eastern Butterflies. Houghton Mifflin Co., Boston, Massachusetts. 396 pp., 48 pls.

Opler, P.A. 1998. A Field Guide to Eastern Butterflies. Houghton Mifflin Co., Boston, Massachusetts. 486 pp., 39 pls.

Opler, P.A. 1999. A Field Guide to Western Butterflies. Houghton Mifflin Co., Boston, Massachusetts. 540 pp., 44 pls.

Opler, P.A. 2003. Fixation of type locality for *Lycaena acmon* Westwood and characterization of the species and its distribution. The Taxonomic Report 4 (1): 1-6.

Opler, P.A. & G.O. Krizek. 1984. Butterflies East of the Great Plains. Johns Hopkins University Press, Baltimore, Maryland. xvii + 294 pp., 54 pls.

Opler, P.A. & J.A. Powell. 1962. Taxonomic and distributional studies on the western components of the *Apodemia mormo* complex (Riodinidae). Journal of the Lepidopterists' Society 15(3):145-171.

Opler, P.A., R.E Stanford & H. Pavulaan, coordinators and editors. 2000. Butterflies of North America. Northern Prairie Wildlife Research Center, Jamestown, North Dakota. [updated periodically]
(<http://www.npwrc.usgs.gov/resource/distr/lepid/bflyusa>)
(also see <http://www.nearctica.com>)

Pavulaan, H. 1995. How many species of azures in Massachusetts, or will the real spring azure please open your wings? Massachusetts Butterflies 6(1):2-4.

Pavulaan, H. & R.R. Gatrell. 1999. A new subspecies of *Brephidium isophthalma* (Lycaenidae: Polyommatinae) from coastal South Carolina. The Taxonomic Report 1(7):1-4.

Pavulaan, H. & D.M. Wright. 2002. *Pterourus appalachensis* (Papilionidae, Papilioninae), a new swallowtail butterfly from the Appalachian region of the United States. The Taxonomic Report 3(7):1-20.

Pavulaan, H. & D.M. Wright. 2005. *Celastrina serotina* (Lycaenidae: Polyommatinae): a new butterfly species from northeastern United States and eastern Canada. The Taxonomic Report 6(6): 1-20.

Penz, C.M. & P.J. DeVries. 2002. Phylogenetic analysis of *Morpho* butterflies (Nymphalidae, Morphinae): implications for classification and natural history. American Museum Novitates 3374:1-33.

Pollock, D.D., W.B. Watt, V.K. Rashbrook, & E.V. Iyengar. 1998. Molecular phylogeny for *Colias* butterflies and their relatives (Lepidoptera: Pieridae). Annals of the Entomological Society of America 91(5):524-531.

Porter, A.H. 1990. Testing nominal species boundaries using gene flow statistics: the taxonomy of two hybridizing admiral butterflies (*Limenitis*: Nymphalidae). *Systematic Zoology* 39(2): 131-147.

Porter, A.H. & H. Geiger. 1988. Genetic and phenotypic population structure of the *Coenonympha tullia* complex (Lepidoptera: Nymphalidae: Satyrinae) in California: no evidence for species boundaries. *Canadian Journal of Zoology* 66(12):2751-2765.

Porter, A.H. & S.O. Mattoon. 1989. A new subspecies of *Coenonympha tullia* (Müller) (Nymphalidae: Satyrinae) confined to the coastal dunes of northern California. *Journal of the Lepidopterists' Society* 43(3):229-238.

Porter, A.H. & A.M. Shapiro. 1991 (1989). Genetics and biogeography of the *Oeneis chryxus* complex (Satyridae) in California. *Journal of Research on the Lepidoptera* 28(4):263-276.

Powell, J.A. 1975. Family Riodinidae. Pages 259-272 in: Howe, W.H., editor. *Butterflies of North America*. Doubleday and Co., Garden City, New York. xiii + 633 pp., 97 pls.

Pratt, G.F. 1988. The evolution and biology of *Euphilotes* biotypes. Ph.D. dissertation, University of California, Riverside. 653 pp.

Pratt, G.F. 1994. Evolution of *Euphilotes* (Lepidoptera, Lycaenidae) by seasonal and host shifts. *Biological Journal of the Linnean Society* 51(1):387-416.

Pratt, G.F. 1999. Battoides group of *Euphilotes*. pp. 236-239 in: Opler, P.A. *A Field Guide to Western Butterflies*. Houghton Mifflin Co., Boston. 540 pp., 44 pls.

Pratt, G.F. & G.R. Ballmer. 1991. Three biotypes of *Apodemia mormo* (Riodinidae) in the Mojave Desert. *Journal of the Lepidopterists' Society* 45(1):46-57.

Pratt, G.F. & G.R. Ballmer. 1993. Correlations of diapause intensities of *Euphilotes* spp. and *Philotella speciosa* (Lepidoptera: Lycaenidae) to host bloom period and elevation. *Annals of the Entomological Society of America* 86(3):265-272.

Pratt, G.F. & J.F. Emmel. 1998. Revision of the *Euphilotes enoptes* and *E. battoides* complexes (Lepidoptera: Lycaenidae). Pp. 207-270 in: Emmel, T.C., editor. *Systematics of Western North American Butterflies*. Mariposa Press, Gainesville, Florida. 878 pp.

Pratt, G.F., D.M. Wright & G.R. Ballmer. 1993 (1991). Multivariate and phylogenetic analyses of larval and adult characters of the Editha complex of the genus *Lycaena* (Lepidoptera: Lycaenidae). *Journal of Research on the Lepidoptera* 30(3/4):175-195.

Pratt, G.F., D.M. Wright & H. Pavulaan. 1994. The various taxa and hosts of the North American *Celastrina* (Lepidoptera: Lycaenidae). *Proceedings of the Entomological Society of Washington* 96(3):566-578.

Pyle, R.M. 1981. The Audubon Society Field Guide to North American Butterflies. Chanticleer Press, Inc., New York. 916 pp.

Pyle, R.M. 1986. The Audubon Society Field Guide to North American Butterflies. Chanticleer Press, Inc., New York. 924 pp.

Pyle, R.M. 2002. Butterflies of Cascadia. Seattle Audubon Society, Seattle, Washington. 420 pp.

Rehn, J.A.G. 1954. The John Eatton Leconte collection of paintings of insects, arachnids, and myriapods. Proceedings of the American Philosophical Society 98(6):442-448.

Riley, N.D. 1975. A Field Guide to the Butterflies of the West Indies. Demeter Press, Inc., Boston. 224 pp., 24 pls.

Riotte, J.C.E. & G. Uchida. 1979 (1978). Butterflies of the Hawaiian Islands, according to the stand of late 1976. Journal of Research on the Lepidoptera 17(1):33-39.

Robbins, R.K. 1988a. Comparative morphology of the butterfly foreleg coxa and trochanter (Lepidoptera) and its systematic implications. Proceedings of the Entomological Society of Washington 90(2):133-154.

Robbins, R.K. 1988b. Male foretarsal variation in Lycaenidae and Riodinidae, and the systematic placement of *Styx infernalis* (Lepidoptera). Proceedings of the Entomological Society of Washington 90(3):356-368.

Robbins, R.K. 1990. The *Mitoura spinetorum* complex in New Mexico and the validity of *M. millerorum* (Lycaenidae: Theclinae). Journal of the Lepidopterists' Society 44(2):68-76.

Robbins, R.K. 1991. Evolution, comparative morphology, and identification of the eumaeine butterfly genus *Rekoa* Kaye (Lycaenidae: Theclinae). Smithsonian Contributions in Zoology 498:1-64.

Robbins, R.K. & G. Lamas. 2004. Eumaeini. Pages 118-137 in: Lamas, G., editor. Checklist: Part 4A. Papilioidea – Hesperioidae in: Heppner, J.B., editor. Atlas of Neotropical Lepidoptera. Scientific Publishers, Gainesville, Florida.

Robbins, R.K. & P.M. Henson. 1986. Why *Pieris rapae* is a better name than *Artogeia rapae* (Pieridae). Journal of the Lepidopterists' Society 40(2):79-92.

Robbins, R.K., G. Lamas, O.H.H. Mielke, D.J. Harvey & M. Casagrande. 1996. Taxonomic composition and ecological structure of the species-rich butterfly community at Pakitza, Parque Nacional del Manu, Peru. Pp. 217-252 in: Wilson, D.E. & A. Sandoval, editors. Manu: The Biodiversity of Southeastern Peru. Smithsonian Institution Press, Washington, D.C. 679 pp.

- Robbins, R.K. & S.S. Nicolay. 1999. Taxonomy and nomenclature of *Strymon istapa* and *S. columella* (Lycaenidae: Theclinae). Journal of the Lepidopterists' Society 52(3):318-327.
- Roever, K. 1975. Family Megathymidae. Pp. 411-422 in: Howe, W.H., editor. The Butterflies of North America. Doubleday and Co., Garden City, New York. xiii + 633 pp., 97 pls.
- Roever, K. 1998. Descriptions of new *Agathymus* (Lepidoptera: Megathymidae) from the southwestern United States. Pp. 491-500 in: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida. 878 pp.
- Roos, P. & W. Arnscheid. 1984. Zur Nomenklatur der nearktischen Arten des *Erebia youngi/dabanensis*-Komplexes (Lepidoptera: Satyridae). Beiträge zur Kenntnis der Erebien, XVI. Nota Lepidopterologica 7(2):178-179.
- Roos, P. & H.P. Kimmich. 1983. Eine neue Art der *Erebia alberganus*-Gruppe aus Nordkanada (Lep.: Satyridae). Entomologische Zeitschrift 93(6):69-75.
- Rydon, A.H.B. 1971. The systematics of the Charaxidae (Lepidoptera, Nymphaloidea). Entomologist's Record and Journal of Variation 83(8):219-233, pls. 8a-8b, figs. 31-36; (9):283-287; (10):310-316, pl. 11; (11):336-341; (12):384-388.
- Sattler, K. & W.G. Tremewan. 1984. The Lepidoptera names of Denis and Schiffmüller - a case for stability. Nota Lepidopterologica 7(3):282-285.
- Schwartz, A. 1989. The Butterflies of Hispaniola. University of Florida Press, Gainesville, Florida. xiv + 580 pp., 8 pls.
- Scott, J.A. 1974. Adult behavior and population biology of *Poladryas minuta* (Nymphalidae) and the relationship of the Texas and Colorado populations. Pan Pacific Entomologist 50(1):9-22.
- Scott, J.A. 1975. Clinal intergradation of *Hesperia comma colorado* (Hesperiidae). Journal of the Lepidopterists' Society 29(3):156-161.
- Scott, J. A. 1977 (1976). *Amblyscirtes "erna,"* a form of *Amblyscirtes aenus*. Journal of Research on the Lepidoptera 15(2):92.
- Scott, J.A. 1978. The identity of the Rocky Mountain *Lycaena dorcas-helloides* complex (Lycaenidae). Journal of Research on the Lepidoptera 17(1):40-50.
- Scott, J.A. 1984. A review of *Polygonia progne* (*oreas*) and *P. gracilis* (*zephyrus*) (Nymphalidae), including a new subspecies from the southern Rocky Mountains. Journal of Research on the Lepidoptera 23(3):197-210.
- Scott, J.A. 1986. The Butterflies of North America: A Natural History and Field Guide. Stanford University Press, Stanford, California. 583 pp., 64 pls.

- Scott, J.A. 1994. Biology and systematics of *Phyciodes* (*Phyciodes*). Papilio, New Series 7:1-120.
- Scott, J.A. 1998. New western North American butterflies. Papilio, New Series 11:1-12.
- Scott, J.A., N.G. Kondla, & S.M. Spomer. 1998. *Speyeria hesperis* and *Speyeria atlantis* are distinct species. Papilio, New Series 8:1-31.
- Scott, J.A. & D.M. Wright. 1993. *Celastrina nigra* and its synonym *C. ebenina* (Lepidoptera: Lycaenidae). Journal of Research on the Lepidoptera 30(3-4):257-260.
- Scott, J.A. & D.M. Wright. 1998. A new *Celastrina* from the Front Range of Colorado. Papilio, New Series 9:1-15.
- Scriber, J.M., M.H. Evans & R.C. Lederhouse. 1989. Hybridization of the Mexican tiger swallowtail *Papilio alexiares garcia* with other *glaucus* group species and survival of pupae and hybrid larvae on potential host plants. Journal of Research on the Lepidoptera 27(3/4):222-232.
- Shapiro, A.M. 1974. Butterflies and skippers of New York State. Search 4(3):1-60.
- Shapiro, A.M. 1976. The biological status of Nearctic taxa in the *Pieris protodice-occidentalis* group (Pieridae). Journal of the Lepidopterists' Society 30(4):289-300.
- Shapiro, A.M. 1984. Review. Systematische unterschungen am *Pieris napi-bryoniae*-komplex (Lepidoptera: Pieridae), by Ulf Eitschberger. Journal of the Lepidopterists' Society 38(4):342-327.
- Shapiro, A.M. & H. Geiger. 1989. Electrophoretic comparisons of vicariant *Vanessa*: genetic differentiation between *V. annabella* and *V. carye* (Nymphalidae) since the Great American Interchange. Journal of the Lepidopterists' Society 43(2):81-92.
- Shepard, J.H. 1998. The correct name for the *Boloria chariclea/titania* complex in North America (Lepidoptera: Nymphalidae). Pp. 727-730 in: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida. 878 pp.
- Shepard, J.H., L. Crabo, & J.P. Pelham. 1998. Revision of *Boloria natazhati* (Gibson) (Lepidoptera: Nymphalidae). Pp. 731-736 in: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida. 878 pp.
- Shepard, J.H. & T.R. Manley. 1998. A species revision of the *Parnassius phoebus* complex in North America (Lepidoptera: Papilionidae). Pp. 717-726 in: Emmel, T.C., editor. Systematics of Western North American Butterflies. Mariposa Press, Gainesville, Florida. 878 pp.

- Shields, O. 1975. Studies on North American *Philotes* (Lycaenidae) IV. Taxonomic and biological notes, and new subspecies. Bulletin of the Allyn Museum 28:1-36.
- Shields, O. 1984. A revised, annotated checklist of world Libytheidae. Journal of Research on the Lepidoptera 22(4):264-266.
- Shields, O. 1985. Zoogeography of Libytheidae (snouts or breaks [sic]). Tokurana (Acta Rhopalocera) 9(1):1-58.
- Shields, O. & J.R. Mori. 1979 (1978). Another *Anthocharis lanceolata* x *A. sara* hybrid. Journal of Research on the Lepidoptera 17(1):53-55.
- Shuey, J.A. 1989. The morpho-species concept of *Euphyes dion* with the description of a new species (Hesperiidae). Journal of Research on the Lepidoptera 27(3-4):160-172.
- Shuey, J.A. 1993. Phylogeny and biogeography of *Euphyes* Scudder (Hesperiidae). Journal of the Lepidopterists' Society 47(4):261-278.
- Silberglied, R., A. Aiello & G. Lamas. 1979. Neotropical butterflies of the genus *Anartia*: systematics, life histories, and general biology (Lepidoptera: Nymphalidae). Psyche 86(2-3):219-260.
- Smith, D.S., L.D. Miller & J.Y. Miller. 1994. The Butterflies of the West Indies and South Florida. Oxford University Press, Oxford. viii + 264 pp., 32 pls.
- Sperling, F.A.H. 1987. Evolution of the *Papilio machaon* species group in western Canada (Lepidoptera: Papilionidae). Quaestiones Entomologicae 23(2):198-315.
- Sperling, F.A.H. 1993a. Mitochondrial DNA phylogeny of the *Papilio machaon* species group (Lepidoptera: Papilionidae). Memoirs of the Entomological Society of Canada 165:233-242.
- Sperling, F.A.H. 1993b. Mitochondrial DNA variation and Haldane's rule in the *Papilio glaucus* and *P. troilus* species groups. Heredity 71(3):227-233.
- Sperling, F.A.H. 2003. Butterfly species and molecular phylogenies. Pages 431-458 in: Boggs, C., W. Watt & P. Ehrlich, editors. Butterflies: Evolution and Ecology Taking Flight. University of Chicago Press, Chicago.
- Sperling, F.A.H. & R.G. Harrison. 1994. Mitochondrial DNA variation within and between species of the *Papilio machaon* group of swallowtail butterflies. Evolution 48(2):408-422.
- Stallings, D.B. & J.R. Turner. 1946. Texas Lepidoptera (Rhopalocera: Papilionoidea). Entomological News 57(2):44-49.
- Stanford, R.E. 2002. Millenial Update on Western USA/ Southern Canada/ Northern Mexico

Butterflies and Skippers. Second Edition. Author, Denver, Colorado. [121] pp.

Stanford, R.E. & P.A. Opler. 1993. Atlas of Western USA Butterflies, Including Adjacent Parts of Canada and Mexico. Authors, Denver and Fort Collins, Colorado. x + 275 pp.

Steinbauer, S.R. 1989. Taxonomic notes and descriptions of new taxa in the Neotropical Hesperiidae. Part I. Pyrginae. Bulletin of the Allyn Museum 127:1-70.

Tilden, J.W. 1952. Concerning the identity of *Mitoura nelsoni muiri*. The Lepidopterists' News 6(6-8):95-96.

Tilden, J.W. 1974. Unusual and interesting butterfly records from Texas. Journal of the Lepidopterists' Society 28(1):22-25.

Tilden, J.W. & A.C. Smith. 1986. A Field Guide to Western Butterflies. Houghton Mifflin Co., Boston. xii + 370 pp., 48 pls.

Toliver, M.E., R. Holland & S.J. Cary. 1994. Distribution of Butterflies in New Mexico (Lepidoptera: Hesperioidea and Papilionoidea). Second Edition. Authors, Albuquerque, New Mexico. [xlvi] + 393 pp.

Troubridge, J.T. & D.K. Parshall. 1988. A review of the *Oeneis polixenes* (Fabricius) complex in North America. The Canadian Entomologist 120(7):679-696.

Troubridge, J.T. & K.W. Philip. 1983 (1982). A review of the *Erebia dabanensis* complex (Lepidoptera: Satyridae), with descriptions of two new species. Journal of Research on the Lepidoptera 21(2):107-146.

Troubridge, J.T., K.W. Philip, J.A. Scott & J.H. Shepard. 1982. A new species of *Oeneis* (Satyridae) from the North American arctic. The Canadian Entomologist 114(10):881-889.

Troubridge, J.T. & K.W. Philip. 1990. A new species of *Colias* (Lepidoptera: Pieridae) from arctic Canada. The Canadian Entomologist 122(1-2):15-20.

Troubridge, J.T. & D.M. Wood. 1995. Biology and taxonomic status of *Boloria natazhati* (Gibson)(Nymphalidae). Journal of the Lepidopterists' Society 44(3):180-187.

Turner, T.W. & J.R. Parnell. 1985. The identification of two species of *Junonia* Hübner (Lepidoptera: Nymphalidae): *J. evarete* and *J. genoveva* in Jamaica. Journal of Research on the Lepidoptera 24(2):142-153.

Tuzov, V.K., P.V. Bogdanov, A.L. Devyatkin, L.V. Kaabak, V.A. Korolev, V.S. Murzin, G.D. Samodurov & E.A. Tarasov. 1997. Guide to the Butterflies of Russia and Surrounding Territories (Lepidoptera: Rhopalocera). Volume 1. Hesperiidae, Papilionidae, Pieridae, Satyridae. Pensoft, Sofia, Bulgaria. 480 pp., 79 pls.

Tuzov, V.K., P.V. Bogdanov, A.L. Devyatkin, L.V. Kaabak, V.A. Korolev, V.S. Murzin, G.D. Samodurov & E.A. Tarasov. 2000. Guide to the Butterflies of Russia and Surrounding Territories (Lepidoptera: Rhopalocera). Volume 2. Libytheidae, Danaidae, Nymphalidae, Riodinidae and Lycaenidae. Pensoft, Sofia, Bulgaria. 581 pp., 88 pls.

Tyler, H., K.S. Brown, Jr. & K. Wilson. 1994. Swallowtail Butterflies of the Americas. A Study in Biological Dynamics, Ecological Diversity, Biosystematics, and Conservation. Scientific Publishers, Gainesville, Florida. 376 pp.

Vargas, I.F., A.M. Luis, J.B. Llorente & A.D. Warren. 1996. Butterflies of the State of Jalisco, Mexico. Journal of the Lepidopterists' Society 50(2):97-138.

Verhulst, J.T. 2000. Les *Colias* du Globe. 2 volumes. Goecke and Evers, Keltern, Germany. 308 pp., 183 pls.

Voss, E.G. 1952. On the classification of the Hesperiidae. Annals of the Entomological Society of America 45(2):246-258.

Wahlberg, N. 2002. [Classification of the Nymphalidae]. [updated periodically] (<http://www.zoologi.su.se/research/wahlberg/Nymphalidae/Classification2.htm>)

Wahlberg, N. & S. Nylin. 2003. Morphology versus molecules: resolution of the positions of *Nymphalis*, *Polygonia*, and related genera (Lepidoptera: Nymphalidae). Cladistics 19(2003):213-223.

Wahlberg, N., R. Oliveira & J.A. Scott. 2003. The phylogenetic relationships of *Phyciodes* butterfly species (Lepidoptera: Nymphalidae): complex mtDNA variation and species delimitations. Systematic Entomology 28: 257-273.

Wahlberg, N. & M. Zimmerman. 2000. Pattern of phylogenetic relationships among members of the tribe Melitaeini (Lepidoptera: Nymphalidae) inferred from mtDNA sequences. Cladistics 16(4):347-368.

Warren, A.D. 1996. Phylogenetic revision of the skippers of the *mithridates* species group, and their replacement in *Eantis* Boisduval, 1836 (Lepidoptera: Hesperiidae: Pyrginae). Entomology Department Honors Thesis, Cornell University, Ithaca, New York. 188 pp., [39] pls.

Warren, A.D. 1997. *Urbanus belli* (Hesperiidae: Pyrginae): a new record for the United States. News of the Lepidopterists' Society 39(3):41,60.

Warren, A.D. 2000. Hesperioidea (Lepidoptera). Pp. 535-580 in: Llorente, J.B., E.S. González & N. Papavero, editors. Biodiversidad, Taxonomía y Biogeografía de Artrópodos de México: Hacia una Síntesis de su Conocimiento. Volumen II. Universidad Nacional Autónoma de México, Mexico City. 676 pp.

Warren, A.D. 2002. Superfamilia Hesperioidea *in:* Lepidoptera *in:* Catálogos de Especies Mexicanas. Comisión Nacional para el Conocimiento y uso de la Biodiversidad, México City. (http://www.conabio.gob.mx/informacion/catalogo_autoridades/doctos/electronicas.html) [updated periodically]

Warren, A.D. 2005. Lepidoptera of North America. 6. Butterflies of Oregon: their taxonomy, distribution, and biology. Contributions of the C.P. Gillette Museum of Arthropod Diversity, Fort Collins, Colo., 408 pp.

Warren, A.D., D.J. Hanson, E. Knudson, and C. Bordelon. 2003. *Achlyodes pallida* (Hesperiidae): a new record for the United States. News of the Lepidopterists' Society 45(4): 128-129.

Warren, A. D. & O. H. H. Mielke. 2005. Synonymic notes on North and Central American *Lerodea* Scudder, 1872 and *Corticea* Evans, 1955 (Lepidoptera, Hesperiidae, Hesperiinae). Revista Brasileira de Zoologia 22(1):285-291.

Warren, A.D., I.F. Vargas, A.M. Luis & J.B. Llorente. 1998. Butterflies of the State of Colima, Mexico. Journal of the Lepidopterists' Society 52(1):40-72.

Warren, B.C.S. 1926. Monograph of the tribe Hesperiidi (European species) with revised classification of the subfamily Hesperiinae (Palaearctic species) based on the genital armature of the males. Transactions of the Entomological Society of London 74(1):1-170.

Warren, B.C.S. 1935. Supplementary notes on the structural characteristics of *Pyrgus freija* (Warren, 1924) (Lepidoptera, Hesperiidae). Transactions of the Royal Entomological Society of London 83(1):127-129.

Warren, B.C.S. 1944. Review of the classification of the Argynnidi: with a systematic revision of the genus *Boloria* (Lepidoptera, Nymphalidae). Transactions of the Royal Entomological Society of London 94(1):1-101, 46 pls.

Warren, B.C.S. 1951. On *Pyrgus freija* (Warren) (Lepidoptera, Hesperiidae). The Entomologist 84(10):217-220, pl. iv.

Warren, B.C.S. 1968. Some structural characteristics of *Erebia inuitica* Wyatt. Entomologists' Record 80(10):242-243.

Watts, J.R. & D.H. Habeck. 1991. Immature stages of *Anthanassa texana seminole* (Lepidoptera: Nymphalidae). Tropical Lepidoptera 2(2):103-106.

Webster, R.P. 1999 (1998). The life history of the maritime ringlet, *Coenonympha tullia nipisiquit* (Satyridae). Journal of the Lepidopterists' Society 52(4):345-355.

Weidenhoffer, Z., G.C. Bozano, & S. Churkin. 2004. Guide to the butterflies of the Palaearctic Region. Lycaenidae, Part II. Subfamily Theclinae, Tribe Eumaeini (partim). Omnes Artes, Milan, Italy, 94 pp.

Wheeler, Q.D. & R. Meier. 2000. Species Concepts and Phylogenetic Theory, a Debate. Columbia University Press, New York. 230 pp.

Wright, D.M. 1995. The American azures: our blue heaven. American Butterflies 3(1):20-28, 30.

Wright, D.M. & H. Pavulaan. 1999. *Celastrina idella* (Lycaenidae: Polyommatinae): a new butterfly species from the Atlantic coastal plain. The Taxonomic Report 1(9):1-11.

Wyatt, C.W. 1958. Observations on *Boloria distincta* (Nymphalidae). The Lepidopterists' News 11(4-5):142-146.

Wyatt, C.W. 1966. Eine neue *Erebia*-art aus Alaska. Zeitschrift der Wiener Entomologischen Gesellschaft 51[77](7):93-94.

Zimmerman, E.C. 1958. Insects of Hawaii. Volume 7. Macrolepidoptera. University of Hawaii Press, Honolulu, ix+542 pp.

Zimmerman, M., N. Wahlberg & H. Decimon. 2000. Phylogeny of *Euphydryas* checkerspot butterflies (Lepidoptera: Nymphalidae) based on mitochondrial DNA sequence data. Annals of the Entomological Society of America 93(3):347-355.

Zirlin, H. 2002. Strangers in a strange land. American Butterflies 10(1):4-11.

Zirlin, H. & J. Ingraham. 1997. Not rotten to the core: butterflies of the Big Apple. American Butterflies 5(3):4-11.

Acknowledgments

We thank the many people who have contributed information on specific records, scientific names, publications, species concepts, and who have provided us with important literature. Notable among them are A. Atkins, G.T. Austin, R. Bailowitz, C.W. Bordelon, Jr., J.P. Brock, E. Brockmann, A.V.Z. Brower, F.M. Brown, J.M. Burns, S.J. Cary, F.S. Chew, H. Chiba, R. de Jong, A.L. Devyatkin, R. Dirig, U. Eitschberger, J.F. Emmel, T.C. Emmel, C.D. Ferris, H.A. Freeman, A.V.L. Freitas, J. Glassberg, P. Hall, P.C. Hammond, R. Hirzel, R. Holland, R. Huber, K. Johnson, A.Y. Kawahara, R.O. Kendall, E. Knudson, N.G. Kondla, J.D. Lafontaine, I. Leeuw, A. Luis-Martínez, J.E. Llorente-Bousquets, C.D. MacNeill, B. Mather, D.V. McCorkle, O.H.H. Mielke, J.S. Miller, J.Y. Miller, L.D. Miller, D.D. Mullins, C.C. Nice, B.A. O'Hara, T.W. Ortenburger, R.S. Peigler, J.A. Powell, G.F. Pratt, F.W & J.D. Preston, R.M. Pyle, M. Quinn, R.K. Robbins, K. Roever, R.L. Romeyn, D.F. Schweitzer, J.A. Scott, A.M. Shapiro, J.H. Shepard, J.A. Shuey, J. Slotten, F.A.H. Sperling, R.E. Stanford, S.R. Steinhauser, S.A. Upson, I. Vargas-Fernández, E.G. Voss, N. Wahlberg, W.B. Watt, D.J. & S.J. Warren, Q.D. Wheeler, D.M. Wright, and especially G. Lamas and J.P. Pelham who assisted in many ways. We are appreciative of their input, but they do not bear responsibility for the final result.